



Degenhardt, L., Grebely, J., Stone, J., Hickman, M., Vickerman, P., Marshall, B. D. L., Bruneau, J., Altice, F. L., Henderson, G., Rahimi-Movaghar, A., & Larney, S. (2019). Global patterns of opioid use and dependence: harms to populations, interventions, and future action. *Lancet*, 394(10208), 1560-1579. [https://doi.org/10.1016/S0140-6736\(19\)32229-9](https://doi.org/10.1016/S0140-6736(19)32229-9)

Peer reviewed version

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1016/S0140-6736\(19\)32229-9](https://doi.org/10.1016/S0140-6736(19)32229-9)

[Link to publication record in Explore Bristol Research](#)
PDF-document

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Webappendices for:

“Global patterns of opioid use and dependence: Population harms, interventions, and future action”

Professor Louisa Degenhardt¹, PhD, Professor Jason Grebely², PhD, Jack Stone³, PhD, Professor Matthew Hickman³, PhD, Professor Peter Vickerman³, PhD, Brandon D.L. Marshall⁴, PhD, Professor Julie Bruneau^{5,6}, MD, Professor Frederick L. Altice⁷, MD, Professor Graeme Henderson⁸, PhD, Professor Afarin Rahimi-Movaghar⁹, PhD, Sarah Larney¹, PhD

Contents

Appendix A: Summary of overall approach to reviewing literature for this paper	5
Panel A1: Types of opioids	5
Web appendix B: Evidence on medicinal uses of opioids.....	6
Panel B1: Medicinal uses of opioids.....	6
References.....	8
Web appendix C: Details of process for reporting and processing of data on opioid analgesic consumption reported to the International Narcotics Control Board	13
Web Appendix D: Data on the prevalence of opioid use collected by the United Nations Office on Drugs and Crime	14
Figure D1: Past-year prevalence of opiate use (top) and extra-medical use of pharmaceutical opioids (bottom)	15
Appendix E: Summary of review of extent of pharmaceutical opioid use and dependence ..	16
Web appendix F: Details of Global Burden of Disease 2017 study data for opioid dependence and opioid overdose	17
Appendix Table F1: Regional and global estimates of the prevalence and number of cases of opioid dependence estimated in the Global Burden of Disease study 2017	19
Web appendix G: Literature on mechanisms and consequences of opioid overdose	20
Search strategy.....	20
Mechanisms of opioid overdose	20
References.....	21
Web appendix H: Details of literature search on other harms elevated among people who use opioids extra-medically.....	22
Review of reviews	22

Injecting risk behaviour.....	22
Dependence	22
HIV incidence	23
HIV injecting risk	23
HCV incidence	23
HCV injecting risk	24
Skin and soft tissue infections	25
Quality of life.....	25
Contact with criminal justice system	25
Accidental injuries.....	26
Suicide	26
Overdose	27
Neonatal outcomes.....	27
Description of evidence base on other harms elevated among people who use opioids extra-medically.....	30
Injecting risk behaviour.....	30
Dependence	30
HIV incidence	30
HIV injecting risk	30
HCV incidence	30
HCV injecting risk	30
Skin and soft tissue infections	30
Quality of life.....	30
Criminal activity	30
Contact with criminal justice system	31
Accidental injuries.....	32
Suicide	32
Overdose	32
Neonatal outcomes.....	32
Web appendix I: Details of literature search on mortality among people who use opioids, including fatal overdose.....	33
Description of methods.....	33

Search strings for electronic literature searches	36
Number of articles identified from opioid mortality search.....	37
PRISMA Flowchart of search strategy	38
List of included studies.....	39
Web appendix J: Details of literature search on reviews of interventions to reduce adverse health outcomes	45
Description of Method	45
Search Strategy	46
Cochrane Library of Systematic Reviews and EBM reviews (Full Text-Cochrane DSR, ACP Journal Club and DARE) search for Level 1 studies.....	46
Peer-reviewed literature database searches for Level 1 studies	47
Description of evidence on reviews of interventions to reduce health outcomes	64
Opioid Agonist Treatment (OAT)	64
Panel J1 – Other opioid agonist treatments (OAT) for opioid dependence	74
Provision of sterile injecting equipment.....	76
Naloxone	77
Drug consumption rooms (DCRs)	78
Oral opioid antagonists.....	79
Sustained release opioid antagonists	81
Residential rehabilitations	82
HIV testing and informing of serostatus.....	82
HCV testing and informing of serostatus.....	82
HIV treatment	83
HCV treatment	83
Suicide prevention strategies	83
Limits on opioid prescribing.....	84
Prescription opioid monitoring programmes	84
Abuse-deterrent opioids.....	85
Compulsory drug detention centres.....	85
Criminalisation of drug use	85
Table J5: Current evidence for effects of interventions to address key outcomes and behaviours among people who use opioids extra-medically	86

Web appendix K: Details of literature search on cost-effectiveness studies to reduce overdose	92
Appendix L: Details of review on retention in treatment, dose of OST and other treatment characteristics	93
Figure L1: Number of people in opioid agonist treatment (OAT) per 100 people who inject drugs, 2015.....	96
Figure L2: Retention in OAT observed in large-scale cohorts	97
Appendix M: Details of mathematical modelling	98
Model Description.....	98
Model Parameterisation and Calibration.....	105
Model Equations	124
Model Outcomes.....	136
References	138

Appendix A: Summary of overall approach to reviewing literature for this paper

This section summarises the broad approach used in searching for evidence during the writing of this paper.

- We systematically searched for high-quality reviews that involved quantitative synthesis where appropriate, in preference to other kinds of reviews. Details of the search strategies used for each component of the paper are summarised in this appendix.
- If no such review could be located, we selected narrative reviews if they clearly described the search strategy and inclusion/exclusion criteria for their review.
- If no such reviews could be located, we conducted searches for high quality single studies with the highest possible quality level for the issue being examined (e.g. well-conducted randomised controlled trials if searching for evidence on an intervention's efficacy).
- If no such studies could be located, we included lower-quality evidence but noted this clearly in tables and text.
- For some key areas, e.g. overdose, only older reviews could be located ^(1,2), so we conducted new full systematic reviews to update and expand upon previous reviews.

Panel A1: Types of opioids

Opioids are substances that stimulate opioid receptors in the body and brain. They can relieve pain, produce euphoria and in high doses, cause respiratory depression, loss of consciousness, overdose and sometimes death. A range of naturally occurring, semi-synthetic and synthetic opioids are produced and are used for medicinal and non-medicinal purposes. Opioids can either partially or fully stimulate opioid receptors, resulting in differing pharmacological responses.

- **Natural opioids (also known as opiates):** Derivatives of naturally occurring alkaloids found in the opium poppy, including morphine, codeine, and opium;
- **Semi-synthetic opioids:** Substances synthesised from naturally occurring opioids, including oxycodone, hydrocodone, hydromorphone, oxymorphone, and heroin (i.e., diacetylmorphine);
- **Synthetic opioids:** Class of drugs designed to mimic naturally occurring opioids, including methadone, fentanyl, pethidine (meperidine), levorphanol, tramadol, tapentadol and dextropropoxyphene.
- **Synthetic opioids considered novel psychoactive substances (NPS),** which particularly include NPS fentanyl analogues such as acetylfentanyl, acryloylfentanyl, butyrylfentanyl, carfentanil, furanylfentanyl, ocfentanyl, methoxyacetylfentanyl, and tetrahydrofuranylfentanyl.

Web appendix B: Evidence on medicinal uses of opioids

The WHO's Model list of essential medicines³ was searched to obtain details of the opioid recommended for medicinal use. It lists codeine, fentanyl, morphine and methadone as opioid analgesics for pain and palliative care. Methadone and buprenorphine are listed for use in the treatment of opioid dependence.

We searched the Cochrane library for reviews of evidence for the use of opioids in treatment of pain, palliative care and opioid dependence. This resulted in 84 reviews that were considered for inclusion. These are listed below. A subset of these were used in the below panel to highlight examples of the nature of evidence for opioids in these indications; the full list of papers is below.

Panel B1: Medicinal uses of opioids

The WHO's Model List of Essential Medicines³ lists codeine, fentanyl, morphine and methadone as essential medicines for pain and palliative care, and methadone and buprenorphine as essential medicines for use in the treatment of opioid dependence. The reference list below contains details of the reviews of evidence for opioids identified in Cochrane, some of which are mentioned below.

Acute pain: Opioids may be used in acute pain conditions including postoperative pain, pain management in labour⁴, and conditions such as acute pancreatitis⁵ and abdominal pain⁶. Low dose and usually less potent opioids are typically used in acute pain (e.g. codeine⁷, 5 or 10 mg oxycodone⁸). There is increasing attention being given to the fact that long-term opioid prescribing can arise out of initial prescribing for an acute pain condition⁹.

Cancer pain: Pain is a common occurrence in people living with cancer and often increases as cancers become more advanced. A recent Cochrane overview concluded that although the amount and quality of evidence on opioids for cancer pain was low, perhaps 19 out of 20 people with moderate to severe cancer pain would derive significant benefit within 14 days; between 10-20% of people would experience intolerable adverse events and change treatments¹⁰. A range of opioids are used in cancer pain; first-line being morphine¹¹, fentanyl¹², oxycodone¹³ and methadone¹⁴. Limited evidence suggests hydromorphone¹⁵ and tapentadol¹⁶ may have similar outcomes compared to morphine or oxycodone. One review concluded buprenorphine should not be considered a first-line opioid for cancer pain¹⁷; another concluded tramadol may be less effective than morphine¹⁸.

Chronic non-cancer pain (CNCP): There is limited, typically low-quality evidence from short-term randomised-controlled trials (RCTs) that opioids in adults living with CNCP may provide

some benefit^{19,20}, but little long-term evaluation²¹. Reviews of specific opioids in specific CNCP conditions such as neuropathic pain have concluded there is limited evidence for some (e.g. oxycodone²²) and insufficient evidence for others (e.g. fentanyl²³, hydromorphone²⁴, buprenorphine²⁵). No RCTs have been undertaken in young people living with CNCP²⁶. There is a lack of evidence of efficacy of high dose (200mg oral morphine equivalent (OME) per day) for CNCP, even in the short term²⁷. The 2016 US CDC guidelines⁹ recommend that clinicians should avoid prescribing opioids, instead prescribing non-pharmacological interventions or non-opioid medications. They placed focus on whether opioids are indicated, careful opioid selection, low dose (<90mg OME/day) and limited prescribing duration.

Opioid withdrawal: using methadone²⁸ or buprenorphine²⁹ in a tapered fashion to manage withdrawal in opioid dependent adults increases likelihood of completing withdrawal (although most people will relapse to opioid use²⁸); it is also associated with overdose and death after completion if people return to use. Completion of withdrawal is improved when psychosocial interventions are also delivered³⁰. Limited evidence suggests opioids used to manage neonatal abstinence syndrome (NAS) reduce time to increased birthweight, but also increase length of hospital stay³¹.

Opioid agonist treatment (OAT): Table 2 in the paper outlines evidence for OAT impacts, which occur across multiple domains. Higher doses of both methadone³² and buprenorphine³³ have been found to improve retention in treatment and reduce extra-medical opioid use (among other outcomes); low-dose buprenorphine has poorer retention than high-dose methadone³³. For those who do not respond to methadone or buprenorphine, diacetylmorphine (i.e., pharmaceutical heroin) increases retention and improves health and social outcomes³⁴ (see **Panel E** for discussion of injectable OAT).

References

1. Ahmad G, O'Flynn H, Attarbashi S, Duffy JM, Watson A. Pain relief for outpatient hysteroscopy. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2010.
2. Amato L, Davoli M, Minozzi S, Ferroni E, Ali R, Ferri M. Methadone at tapered doses for the management of opioid withdrawal. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2013.
3. Amato L, Minozzi S, Davoli M, Vecchi S. Psychosocial and pharmacological treatments versus pharmacological treatments for opioid detoxification. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2011.
4. Amato L, Minozzi S, Davoli M, Vecchi S. Psychosocial combined with agonist maintenance treatments versus agonist maintenance treatments alone for treatment of opioid dependence. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2011.
5. Bao YJ, Hou W, Kong XY, Yang L, Xia J, Hua BJ, et al. Hydromorphone for cancer pain. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2016.
6. Barnes H, McDonald J, Smallwood N, Manser R. Opioids for the palliation of refractory breathlessness in adults with advanced disease and terminal illness. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2016.
7. Basurto OX, Rigau CD, Urrútia G. Opioids for acute pancreatitis pain. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2013.
8. Bellù R, de WKA, Zanini R. Opioids for neonates receiving mechanical ventilation. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2008.
9. Brown R, Howard R, Candy B, Sampson EL. Opioids for agitation in dementia. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2015.
10. Cepeda MS, Camargo F, Zea C, Valencia L. Tramadol for osteoarthritis. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2006.
11. Chaparro LE, Furlan AD, Deshpande A, Mailis-Gagnon A, Atlas S, Turk DC. Opioids compared to placebo or other treatments for chronic low-back pain. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2013.
12. Chaparro LE, Wiffen PJ, Moore RA, Gilron I. Combination pharmacotherapy for the treatment of neuropathic pain in adults. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2012.
13. Cooper TE, Chen J, Wiffen PJ, Derry S, Carr DB, Aldington D, et al. Morphine for chronic neuropathic pain in adults. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2017.
14. Cooper TE, Fisher E, Gray AL, Krane E, Sethna N, van TMA, et al. Opioids for chronic non-cancer pain in children and adolescents. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2017.
15. da CBR, Nüesch E, Kasteler R, Husni E, Welch V, Rutjes AW, et al. Oral or transdermal opioids for osteoarthritis of the knee or hip. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2014.
16. de OCO, Carvalho LB, Carlos K, Conti C, de OMM, Prado LB, et al. Opioids for restless legs syndrome. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2016.
17. Derry S, Derry CJ, Moore RA. Single dose oral ibuprofen plus oxycodone for acute postoperative pain in adults. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2013.

18. Derry S, Moore RA, McQuay HJ. Single dose oral codeine, as a single agent, for acute postoperative pain in adults. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2010.
19. Derry S, Stannard C, Cole P, Wiffen PJ, Knaggs R, Aldington D, et al. Fentanyl for neuropathic pain in adults. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2016.
20. Duehmke RM, Derry S, Wiffen PJ, Bell RF, Aldington D, Moore RA. Tramadol for neuropathic pain in adults. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2017.
21. Eccleston C, Fisher E, Thomas KH, Hearn L, Derry S, Stannard C, et al. Interventions for the reduction of prescribed opioid use in chronic non-cancer pain. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2017.
22. Els C, Jackson TD, Hagtvedt R, Kunyk D, Sonnenberg B, Lappi VG, et al. High-dose opioids for chronic non-cancer pain: an overview of Cochrane Reviews. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2017.
23. Els C, Jackson TD, Kunyk D, Lappi VG, Sonnenberg B, Hagtvedt R, et al. Adverse events associated with medium- and long-term use of opioids for chronic non-cancer pain: an overview of Cochrane Reviews. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2017.
24. Faggiano F, Vigna-Taglianti F, Versino E, Lemma P. Methadone maintenance at different dosages for opioid dependence. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2003.
25. Ferri M, Davoli M, Perucci CA. Heroin maintenance for chronic heroin-dependent individuals. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2011.
26. Ferri M, Minozzi S, Bo A, Amato L. Slow-release oral morphine as maintenance therapy for opioid dependence. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2013.
27. Gardiner SJ, Chang AB, Marchant JM, Petsky HL. Codeine versus placebo for chronic cough in children. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2016.
28. Gaskell H, Derry S, Moore RA, McQuay HJ. Single dose oral oxycodone and oxycodone plus paracetamol (acetaminophen) for acute postoperative pain in adults. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2009.
29. Gaskell H, Derry S, Stannard C, Moore RA. Oxycodone for neuropathic pain in adults. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2016.
30. Gaskell H, Moore RA, Derry S, Stannard C. Oxycodone for pain in fibromyalgia in adults. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2016.
31. Gowing L, Ali R, White JM, Mbewe D. Buprenorphine for managing opioid withdrawal. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2017.
32. Gowing L, Farrell MF, Bornemann R, Sullivan LE, Ali R. Oral substitution treatment of injecting opioid users for prevention of HIV infection. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2011.
33. Guay J, Kopp S. Epidural pain relief versus systemic opioid-based pain relief for abdominal aortic surgery. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2016.
34. Guay J, Nishimori M, Kopp S. Epidural local anaesthetics versus opioid-based analgesic regimens for postoperative gastrointestinal paralysis, vomiting and pain after abdominal surgery. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2016.
35. Gurusamy KS, Vaughan J, Toon CD, Davidson BR. Pharmacological interventions for prevention or treatment of postoperative pain in people undergoing laparoscopic cholecystectomy. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2014.

36. Hadley G, Derry S, Moore RA, Wiffen PJ. Transdermal fentanyl for cancer pain. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2013.
37. Haroutounian S, McNicol ED, Lipman AG. Methadone for chronic non-cancer pain in adults. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2012.
38. Jones L, Othman M, Dowswell T, Alfirevic Z, Gates S, Newburn M, et al. Pain management for women in labour: an overview of systematic reviews. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2012.
39. Jones, A.A., Vila-Rodriguez, F., Leonova, O., Langheimer, V., Lang, D.J., Barr, A.M., Procyshyn, R.M., ... & Honer, W.G. (2015). Mortality from treatable illnesses in marginally housed adults: A prospective cohort study. *BMJ Open*, 5(8), e008876. doi: 10.1136/bmjopen-2015-008876
40. Manterola C, Vial M, Moraga J, Astudillo P. Analgesia in patients with acute abdominal pain. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2011.
41. Marks JL, Colebatch AN, Buchbinder R, Edwards CJ. Pain management for rheumatoid arthritis and cardiovascular or renal comorbidity. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2011.
42. Mason M, Cates CJ, Smith I. Effects of opioid, hypnotic and sedating medications on sleep-disordered breathing in adults with obstructive sleep apnoea. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2015.
43. Mattick RP, Breen C, Kimber J, Davoli M. Methadone maintenance therapy versus no opioid replacement therapy for opioid dependence. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2009.
44. Mattick RP, Breen C, Kimber J, Davoli M. Buprenorphine maintenance versus placebo or methadone maintenance for opioid dependence. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2014.
45. McNicol ED, Ferguson MC, Hudcova J. Patient controlled opioid analgesia versus non-patient controlled opioid analgesia for postoperative pain. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2015.
46. McNicol ED, Ferguson MC, Schumann R. Methadone for neuropathic pain in adults. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2017.
47. McNicol ED, Midbari A, Eisenberg E. Opioids for neuropathic pain. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2013.
48. Minozzi S, Amato L, Bellisario C, Davoli M. Maintenance treatments for opiate -dependent adolescents. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2014.
49. Minozzi S, Amato L, Bellisario C, Ferri M, Davoli M. Maintenance agonist treatments for opiate-dependent pregnant women. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2013.
50. Mkontwana N, Novikova N. Oral analgesia for relieving post-caesarean pain. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2015.
51. Moore RA, Derry S, Aldington D, Wiffen PJ. Adverse events associated with single dose oral analgesics for acute postoperative pain in adults - an overview of Cochrane reviews. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2015.
52. Moore RA, Edwards J, Derry S, McQuay HJ. Single dose oral dihydrocodeine for acute postoperative pain. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2000.
53. Murphy A, O'Sullivan R, Wakai A, Grant TS, Barrett MJ, Cronin J, et al. Intranasal fentanyl for the management of acute pain in children. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2014.

54. Nicholson AB, Watson GR, Derry S, Wiffen PJ. Methadone for cancer pain. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2017.
55. Nielsen S, Larance B, Degenhardt L, Gowing L, Kehler C, Lintzeris N. Opioid agonist treatment for pharmaceutical opioid dependent people. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2016.
56. Noble M, Treadwell JR, Tregear SJ, Coates VH, Wiffen PJ, Akafo C, et al. Long-term opioid management for chronic noncancer pain. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2010.
57. Osborn DA, Jeffery HE, Cole MJ. Opiate treatment for opiate withdrawal in newborn infants. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2010.
58. Quigley C. Opioid switching to improve pain relief and drug tolerability. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2013.
59. Radner H, Ramiro S, Buchbinder R, Landewé RB, van dHD, Aletaha D. Pain management for inflammatory arthritis (rheumatoid arthritis, psoriatic arthritis, ankylosing spondylitis and other spondyloarthritis) and gastrointestinal or liver comorbidity. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2012.
60. Rahimi-Movaghar A, Amin-Esmaeili M, Hefazi M, Yousefi-Nooraie R. Pharmacological therapies for maintenance treatments of opium dependence. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2013.
61. Ramiro S, Radner H, van dHD, van TA, Buchbinder R, Aletaha D, et al. Combination therapy for pain management in inflammatory arthritis (rheumatoid arthritis, ankylosing spondylitis, psoriatic arthritis, other spondyloarthritis). Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2011.
62. Renner R-M, Jensen JTJ, Nichols MDN, Edelman A. Pain control in first trimester surgical abortion. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2009.
63. Santos J, Alarcão J, Fareleira F, Vaz-Carneiro A, Costa J. Tapentadol for chronic musculoskeletal pain in adults. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2015.
64. Saulle R, Vecchi S, Gowing L. Supervised dosing with a long-acting opioid medication in the management of opioid dependence. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2017.
65. Schmidt-Hansen M, Bennett MI, Arnold S, Bromham N, Hilgart JS. Oxycodone for cancer-related pain. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2017.
66. Schmidt-Hansen M, Bromham N, Taubert M, Arnold S, Hilgart JS. Buprenorphine for treating cancer pain. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2015.
67. Schnabel A, Reichl SU, Meyer-Frießem C, Zahn PK, Pogatzki-Zahn E. Tramadol for postoperative pain treatment in children. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2015.
68. Schnabel A, Reichl SU, Zahn PK, Pogatzki-Zahn E. Nalbuphine for postoperative pain treatment in children. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2014.
69. Stannard C, Gaskell H, Derry S, Aldington D, Cole P, Cooper TE, et al. Hydromorphone for neuropathic pain in adults. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2016.
70. Straube C, Derry S, Jackson KC, Wiffen PJ, Bell RF, Strassels S, et al. Codeine, alone and with paracetamol (acetaminophen), for cancer pain. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2014.

71. Toms L, Derry S, Moore RA, McQuay HJ. Single dose oral paracetamol (acetaminophen) with codeine for postoperative pain in adults. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2009.
72. Ullah H, Samad K, Khan FA. Continuous interscalene brachial plexus block versus parenteral analgesia for postoperative pain relief after major shoulder surgery. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2014.
73. Ullman R, Smith LA, Burns E, Mori R, Dowswell T. Parenteral opioids for maternal pain management in labour. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2010.
74. Weibel S, Jelting Y, Afshari A, Pace NL, Eberhart LH, Jokinen J, et al. Patient-controlled analgesia with remifentanyl versus alternative parenteral methods for pain management in labour. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2017.
75. Werawatganon T, Charuluxananan S. Patient controlled intravenous opioid analgesia versus continuous epidural analgesia for pain after intra-abdominal surgery. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2013.
76. Whittle SL, Richards BL, Husni E, Buchbinder R. Opioid therapy for treating rheumatoid arthritis pain. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2011.
77. Wiffen PJ, Derry S, Moore RA. Impact of morphine, fentanyl, oxycodone or codeine on patient consciousness, appetite and thirst when used to treat cancer pain. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2014.
78. Wiffen PJ, Derry S, Moore RA. Tramadol with or without paracetamol (acetaminophen) for cancer pain. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2017.
79. Wiffen PJ, Derry S, Moore RA, Stannard C, Aldington D, Cole P, et al. Buprenorphine for neuropathic pain in adults. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2015.
80. Wiffen PJ, Derry S, Naessens K, Bell RF. Oral tapentadol for cancer pain. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2015.
81. Wiffen PJ, Wee B, Derry S, Bell RF, Moore RA. Opioids for cancer pain - an overview of Cochrane reviews. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2017.
82. Wiffen PJ, Wee B, Moore RA. Oral morphine for cancer pain. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2016.
83. Zeppetella G, Davies AN. Opioids for the management of breakthrough pain in cancer patients. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2015.
84. Zou Z, An MM, Xie Q, Chen XY, Zhang H, Liu GJ, et al. Single dose intra-articular morphine for pain control after knee arthroscopy. Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd; 2016.

Web appendix C: Details of process for reporting and processing of data on opioid analgesic consumption reported to the International Narcotics Control Board

Please see below the description from INCB of the process for collecting and reporting on data on opioid analgesic consumption:

The [1961 Convention](#), which was expanded and strengthened by the 1972 Protocol, consolidated all previous conventions and streamlined the international drug control machinery. The 1961 Convention establishes strict controls on the cultivation of opium poppy, coca bush, cannabis plant and their products, which, in the Convention, are described as "narcotic drugs" (although cocaine is a stimulant drug rather than one that induces sleep). Control is exercised over [124 narcotic drugs](#), mainly natural products, such as opium and its derivatives, morphine, codeine and heroin, but also synthetic drugs, such as methadone and pethidine, as well as cannabis and coca leaf. [Parties to the 1961 Convention](#) undertake to limit the production, manufacture, export, import, distribution and stocks of, trade in and use and possession of the controlled drugs so that they are used exclusively for medical and scientific purposes. The production and distribution of controlled substances must be licensed and supervised, and Governments must provide estimates and statistical returns to INCB [on the forms supplied for that purpose](#), on the quantities of drugs required, manufactured and utilized and the quantities seized by police and customs officers. The control system established under the 1961 Convention functions well, and the system of estimates first introduced by the 1931 Convention is considered to be the key to that success. The system of estimates covers all States, regardless of whether or not they are parties to the 1961 Convention. Each year, INCB publishes in a [technical publication](#) information about the licit movement of the internationally controlled narcotic drugs.

All national competent authorities have to submit to INCB the following forms with the information Governments are required to furnish to the Board pursuant to the provisions of the Single Convention:

[Form A](#) (Quarterly Statistics of Imports and Exports of Narcotic Drugs; to be submitted four times a year);

[Form B](#) (Annual Estimates of Requirements of Narcotic Drugs, Manufacture of Synthetic Drugs, Opium Production and Cultivation of the Opium Poppy for Purposes other than Opium Production); and

[Form C](#) (Annual Statistics of Production, Manufacture, Consumption, Stocks and Seizures of Narcotic Drugs).


If any Government has to modify its estimates, which were furnished to INCB in Form B, the national competent authority shall send to INCB the form entitled [Supplement to Form B](#) (Supplement to the Annual Estimates of Requirements of Narcotic Drugs).

Web Appendix D: Data on the prevalence of opioid use collected by the United Nations Office on Drugs and Crime

Data on the annual prevalence of opioids use as a percentage of the population age 15-64 were obtained from the UNODC World Drug Report 2018, available from <https://www.unodc.org/wdr2017/en/maps-and-graphs.html>. Data from individual countries are mainly sourced from UNODC's Annual Report Questionnaire (ARQ), which collects and reports data on extent, patterns and trends in drug use and its health consequences across the world. The World Drug Report also incorporates data from the literature and grey literature, such as from government sources, academic research, and major international surveys that monitor drug use.

Estimates of the prevalence of opioid use are computed using various adjustments. The figure below presents UNODC's estimates of the prevalence of opiate and extra-medical pharmaceutical opioids in the past year for 2016.

Figure D1: Past-year prevalence of opiate use (top) and extra-medical use of pharmaceutical opioids (bottom)



Source: United Nations Office on Drugs and Crime's 2018 World Drug Report³⁵ data tables.

Appendix E: Summary of review of extent of pharmaceutical opioid use and dependence

Ellerstrand et al (2018)³⁶ conducted a systematic review of the prevalence of the prevalence of illicit and pharmaceutical opioid (PO) nonmedical use and dependence by country were extracted. The search for this study was undertaken as a part of the GBD 2016 study on the global prevalence of substance dependence. Multiple research strategies were utilised, including systematic searches of peer-reviewed literature and collaboration with experts in the field. The choice of databases was selected in concert with expert opinion as well as discussion with university research librarians. A systematic search of peer-reviewed literature was performed using Medline, Embase and PsycINFO databases. The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) style was employed ³⁷.

Search terms were grouped into: 1. substances; 2. use or dependence; 3. database specific terms representing substance dependence (eg. MeSH or Emtree terms); and 4. epidemiology measures (eg. prevalence). Search strategy was grouped as ((1 AND 2) OR 3) AND 4. Epidemiological search strings were standardised for the Psychiatric Epidemiology and Burden of Disease Group (PEABOD team) of which the researchers were a part. This team researches mental and substance use disorders in the GBD study. Search limitations comprise publication year from 2009 onwards (as the last systematic review on opioid dependence was conducted in 2009), peer reviewed journal articles and human studies. There were no limits set on language; the research team included translators. All stages of screening were performed by two reviewers. Data extraction was conducted simultaneously with full-screen review i.e. if an article was selected for inclusion, it was immediately extracted.

Web appendix F: Details of Global Burden of Disease 2017 study data for opioid dependence and opioid overdose

Burden was quantified by geography, for 20 age groups covering 0-99 years, both sexes, and timepoints spanning the period 1990 to 2017. Comprehensive methods for estimating YLDs, YLLs, and DALYs have been presented in a series of publications³⁸⁻⁴¹

YLLs

Input data on causes of death data came from vital registration, verbal autopsy and surveillance databases dating back to 1980.^{38,41} Normative life tables were generated using data on the lowest death rates for each age group within geographies with total populations of more than 5 million. YLLs were then estimated by multiplying cause-specific deaths at a given age by the standard life expectancy at that age obtained from this normative life table. Full details of all the modelling process have been published previously.^{38,41}

The Cause of Death Ensemble modelling (CODEm) strategy was used to model cause of death data by location, age, sex, and year for opioids.^{38,41} CODEm model outputs for all GBD causes were then rescaled to fit the all-cause mortality envelope to derive final cause-specific deaths; deaths coded as alcohol and drug poisonings were attributed to the relevant alcohol and drug use disorders.

YLDs

Epidemiological disease models

Systematic reviews of the literature were conducted to compile data on the prevalence, incidence, remission, and excess mortality associated with opioid dependence. Electronic databases (PubMed, EMBASE, PsycINFO) and grey literature sources were searched in accordance with the Preferred Reporting Items for Systematic Reviews (PRISMA) guidelines.⁴² For each epidemiological parameter, eligible estimates were derived from studies published since 1980.

The epidemiological data obtained from systematic literature reviews were modelled in DisMod-MR 2.1. This is a Bayesian meta-regression tool which pools data-points from different sources and adjusts for known sources of variability (e.g. differences in case definitions and sampling method) to produce internally consistent estimates of incidence, prevalence, remission, and excess mortality.⁴³ It is also used to estimate the epidemiology of disorders within areas with limited data.

DisMod-MR 2.1 analyses ran in a sequence of estimations at each level of the GBD geographic hierarchy (global, super-region, region, country, and if applicable, subnational locations) with consistency imposed between estimates at each level. More information on DisMod-MR 2.1 has been presented in other publications.³⁸⁻⁴¹

Although our inclusion criteria ensured minimum study quality, there was considerable variability between studies that reflected different methodologies and analyses (e.g. see⁴⁴⁻⁴⁸). Uncertainty in both the epidemiological data and in modelling was propagated to the final prevalence output used to calculate YLDs. This was in addition to the uncertainty from fixed effects and random effects for country and regions.⁴³

Disability weights

Each country, age, sex, and year-specific prevalence derived by DisMod MR 2.1 was multiplied by a disorder-specific disability weight to estimate YLDs. Disability weights ranged from 0 ('perfect health') to 1 ('death') to quantify the severity of the health loss associated with a given disease or injury. They were obtained from population surveys in a number of different countries and from an open-access survey available in multiple languages in which lay participants were presented with pairs of short descriptions of health states and asked to rate which they considered the more 'unhealthy'.⁴⁹⁻⁵¹ Disability weights were generated for all sequelae of diseases and injuries. Further details regarding disability weight methodology has been published previously⁵⁰⁻⁵².

Disability-adjusted life-years (DALYs)

The burden calculation involved (1) aggregating substance use disorder-specific epidemiological data and disability weights to calculate prevalent YLDs⁴⁰; (2) multiplying substance use disorder-specific estimates of mortality by standard life expectancy at the age of death to calculate YLLs^{38,41}; and (3) summing YLDs and YLLs to generate substance use disorder-specific DALYs³⁹.

DALYs were derived by summing YLD and YLLs for each disorder, location, age group, sex, and year. Age-standardised rates of prevalence, deaths, YLLs, YLDs, and DALYs were estimated using the GBD world population age standard. Uncertainty was derived for all estimates by simulating 1000 draws from each estimate's posterior distribution. This captures uncertainty arising from primary inputs, sample sizes in the data collected, adjustments made to the data during modelling, and model estimation. For YLLs, they capture uncertainty due to study sample sizes, adjustments made to the all-cause mortality data, and model estimation.

Appendix Table F1: Regional and global estimates of the prevalence and number of cases of opioid dependence estimated in the Global Burden of Disease study 2017

Location	Opioid dependence		Opioid overdose deaths	
	Cases of opioid dependence (1000s) (95%UI)	Age-standardised rate per 100,000 (95%UI)	Number of opioid deaths (1000s) (95%UI)	Age-standardised overdose rate per 100,000 (95%UI)
Andean Latin America	205.79 (169.91 - 248.55)	331.19 (274.4 - 397.30)	0.48 (0.42 - 0.54)	0.84 (0.74 - 0.94)
Australasia	124.07 (105.71 - 144.66)	430.27 (363.31 - 507.77)	0.69 (0.60 - 0.79)	2.19 (1.90 - 2.52)
Caribbean	155.12 (127.32 - 185.88)	319.47 (261.98 - 383.75)	0.31 (0.28 - 0.35)	0.63 (0.56 - 0.70)
Central Asia	202.10 (169.27 - 239.75)	208.65 (176.12-246.45)	1.17 (1.08 - 1.29)	1.30 (1.21 - 1.43)
Central Europe	225.37 (191.84 - 263.90)	179.21 (149.73 - 212.86)	0.73 (0.69 - 0.77)	0.5 (0.47 - 0.53)
Central Latin America	875.53 (724.77 - 1057.10)	330.94 (274.96 - 399.12)	1.71 (1.64 - 1.79)	0.68 (0.65 - 0.71)
Central sub-Saharan Africa	353.71 (286.13 - 437.21)	335.12 (275.11 - 408.19)	0.46 (0.33 - 0.60)	0.53 (0.38 - 0.70)
East Asia	10,501.21(8,745.54 – 12,643.64)	622.21 (509.85 - 759.44)	15.75 (14.81 -16.86)	0.85 (0.80 - 0.91)
Eastern Europe	712.91 (606.90 - 834.42)	307.76 (258.42 - 366.14)	8.26 (7.82 - 8.81)	3.31 (3.13 - 3.54)
Eastern sub-Saharan Africa	988.30 (794.10 – 1,241.43)	294.72 (242.63 - 361.19)	0.68 (0.44 - 0.91)	0.28 (0.17 - 0.37)
High-Income Asia Pacific	497.33 (418.84 - 588.46)	260.54 (212.96 - 317.09)	0.55 (0.53 - 0.58)	0.19 (0.186 - 0.20)
High-Income North America	4,751.86 (4,058.04 – 5,574.96)	1,347.66 (1,136.95 – 1,609.35)	48.63 (45.10 - 52.08)	12.31 (11.46 - 13.14)
North Africa and Middle East	9,286.71 (7,733.90 – 11,004.65)	1,459.36 (1,226.44 – 1,718.77)	8.31 (7.77 - 9.04)	1.46 (1.37 - 1.58)
Oceania	29.78 (23.76 - 37.68)	244.06 (198.40 - 305.06)	0.02 (0.009 - 0.03)	0.18 (0.10 - 0.25)
South Asia	6,331.35 (5,141.18 – 7,769.83)	346.80 (284.75 - 421.59)	8.14 (6.10 - 9.37)	0.53 (0.39 - 0.61)
Southeast Asia	1,330.32 (1,099.55 -1,622.60)	189.16 (156.40 - 230.81)	3.02 (2.77 - 3.28)	0.46 (0.42 - 0.50)
Southern Latin America	183.96 (151.39 - 225.12)	264.69 (215.46 - 326.73)	0.52 (0.47 - 0.58)	0.67 (0.61 - 0.75)
Southern sub-Saharan Africa	426.59 (349.60 - 523.44)	516.09 (428.18 - 625.68)	0.65 (0.58 - 0.78)	0.86 (0.77 - 1.03)
Tropical Latin America	796.39 (964.86 - 656.66)	333.88 (275.23 - 404.91)	1.31 (1.26 - 1.38)	0.56 (0.53 - 0.58)
Western Europe	1,188.14 (1037.55 – 1,343.14)	262.50 (226.55 - 300.27)	5.06 (4.86 - 5.41)	1.02 (0.98 - 1.10)
Western sub-Saharan Africa	1,318.05 (1077.19 – 1,606.51)	356.12 (297.96 - 424.97)	3.05 (2.40 - 3.82)	0.98 (0.80 - 1.25)
Global	40,484.59 (34,271.36 – 47,941.59)	510.28 (430.32 - 605.55)	109.52 (105.75 – 113.56)	1.37 (1.32 - 1.42)

Web appendix G: Literature on mechanisms and consequences of opioid overdose

Search strategy

This section was written following a series of literature searches. These were performed on PubMed and included searches for ‘heroin OR morphine OR opioid’ and each of the following terms – pulmonary function, pulmonary oedema, respiratory depression, brain imaging, brain damage, apoptosis, cognitive function, poly drug use, poly pharmacology, pregabalin, gabapentin – as well as searches for ‘fentanyl’ and respiratory depression, muscle rigidity. Papers were extracted by the authors for their relevance to the topics being reviewed on the basis of their titles and abstracts. In addition, secondary references cited in extracted papers were consulted. Where available recent meta analyses are referenced.

Mechanisms of opioid overdose

At higher doses, opioids suppress respiratory rhythm generation and reduce normal physiological responsiveness of central and peripheral chemoreceptors. As a result, CO₂ levels rise and hypoxia develops¹. Consequently, breathing rates decrease and eventually stop.

Though respiratory depression contributes most to fatal overdose, aspiration of oral or gastric contents may also occur. Time to opioid-related death is variable. Death can occur quickly (e.g. 10-30 minutes when injected), or longer when administered using other routes²⁻⁴. When more prolonged, other pathophysiological responses like respiratory acidosis, pulmonary oedema and aspiration pneumonia increase morbidity or can themselves result in death.

Fentanyl and its structural analogues are potent, highly lipid soluble opioids that rapidly enter the brain following intravenous injection, resulting in accelerated respiratory depression⁵. Though not fully understood⁶, they also induce “respiratory paralysis” by promoting higher levels of chest wall rigidity by causing the respiratory intercostal and diaphragmatic muscles to constrict and prevent chest inflation⁶, relative to other opioids.

For people who use opioids long-term, tolerance to opioids develops, requiring larger doses to either avoid withdrawal symptoms and/or achieve euphoria. Because tolerance to respiratory depression is believed to be less than it is for euphoria (or analgesia in animal models)⁷, users will often increase the dose of opioids enhancing the effects of respiratory depression⁷⁻⁹.

References

1. Pattinson KTS. Opioids and the control of respiration. *British Journal of Anaesthesia* 2008; **100**(6): 747-58.
2. Darke S, Duflou J. The toxicology of heroin-related death: estimating survival times. *Addiction* 2016; **111**(9): 1607-13.
3. Goldberger BA, Cone EJ, Grant TM, Caplan YH, Levine BS, Smialek JE. Disposition of heroin and its metabolites in heroin-related deaths. *J Anal Toxicol* 1994; **18**(1): 22-8.
4. Hickman M, Carrivick S, Paterson S, et al. London audit of drug-related overdose deaths: characteristics and typology, and implications for prevention and monitoring. *Addiction* 2007; **102**(2): 317-23.
5. Suzuki J, El-Haddad S. A review: Fentanyl and non-pharmaceutical fentanyls. *Drug and Alcohol Dependence* 2017; **171**: 107-16.
6. Burns G, DeRienz RT, Baker DD, Casavant M, Spiller HA. Could chest wall rigidity be a factor in rapid death from illicit fentanyl abuse? *Clinical Toxicology* 2016; **54**(5): 420-3.
7. White JM, Irvine RJ. Mechanisms of fatal opioid overdose. *Addiction* 1999; **94**(7): 961-72.
8. Hill R, Dewey WL, Kelly E, Henderson G. Oxycodone-induced tolerance to respiratory depression: reversal by ethanol, pregabalin and protein kinase C inhibition. *Br J Pharmacol* 2018; **175**(12): 2492-503.
9. Hill R, Lyndon A, Withey S, et al. Ethanol Reversal of Tolerance to the Respiratory Depressant Effects of Morphine. *Neuropsychopharmacology* 2016; **41**(3): 762-73.

Web appendix H: Details of literature search on other harms elevated among people who use opioids extra-medically

Review of reviews

Injecting risk behaviour

Medline search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp heroin/ or exp opioid-related disorders/	86559
2	(inject* or injecting risk or (risk behaviour adj3 inject*)).ti,ab.	613357
3	1 and 2	12944
4	limit 3 to (human and yr="2000 -Current" and "review")	409

Embase search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp opiate/ or exp heroin dependence/	151122
2	(inject* or injecting risk or (risk behaviour adj3 inject*)).ti,ab.	989749
3	1 and 2	18031
4	limit 3 to (human and yr="2000 -Current" and "review")	700

Dependence

Medline search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp heroin/ or exp opioid-related disorders/	86559
2	((dependen\$ or use-disorder) and risk).ti,ab.	65579
3	1 and 2	1643
4	limit 3 to (humans and yr="2000 -Current" and "review")	301

Embase search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp opiate/ or exp heroin dependence/	151197
2	((dependen\$ or use-disorder) and risk).ti,ab.	109741
3	1 and 2	3175
4	limit 3 to (human and yr="2000 -Current" and "review")	389

HIV incidence

Medline search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp heroin/ or exp opioid-related disorders/	86559
2	(HIV adj3 incidence).ti,ab or (exp HIV/ and exp Incidence/)	5021
3	1 and 2	119
4	limit 3 to (human and yr="2000 -Current" and "review")	16

Embase search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp opiate/ or exp heroin dependence/	151197
2	(HIV adj3 incidence).ti,ab. or (exp Human Immunodeficiency Virus/ and exp Incidence/)	8223
3	1 and 2	174
4	limit 3 to (human and yr="2000 -Current" and "review")	20

HIV injecting risk

Medline search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp heroin/ or exp opioid-related disorders/	86559
2	(HIV adj3 inject*).ti,ab. or (exp HIV/ and (exp Intravenous Drug Abuse/ or inject*.ti,ab. or injecting risk.ti,ab.))	4365
3	1 and 2	432
4	limit 3 to (human and yr="2000 -Current" and "review")	31

Embase search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp opiate/ or exp heroin dependence/	151197
2	(HIV adj3 inject*).ti,ab. or (exp Human Immunodeficiency Virus/ and (exp Intravenous Drug Abuse/ or inject*.ti,ab. or injecting risk.ti,ab.))	8231
3	1 and 2	875
4	limit 3 to (human and yr="2000 -Current" and "review")	55

HCV incidence

Medline search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp heroin/ or exp opioid-related disorders/	86581
2	((exp Hepatitis C/ or "HCV".ti,ab.) and exp Incidence/) or (HCV adj3 incidence).ti,ab	2672
3	1 and 2	87
4	limit 3 to (human and yr="2000 -Current" and "review")	11

Embase search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp opiate/ or exp heroin dependence/	151197
2	((exp Hepatitis C/ or "HCV".ti,ab.) and exp Incidence/) or (HCV adj3 incidence).ti,ab	5649
3	1 and 2	165
4	limit 3 to (human and yr="2000 -Current" and "review")	21

HCV injecting risk

Medline search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp heroin/ or exp opioid-related disorders/	86581
2	((exp Hepatitis C/ or "HCV".ti,ab.) and (exp Intravenous Drug Abuse/ or inject*.ti,ab. or injecting risk.ti,ab.)) or (HCV adj3 inject*).ti,ab	4672
3	1 and 2	532
4	limit 3 to (human and yr="2000 -Current" and "review")	45

Embase search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp opiate/ or exp heroin dependence/	151197
2	((exp Hepatitis C/ or "HCV".ti,ab.) and (exp Intravenous Drug Abuse/ or inject*.ti,ab. or injecting risk.ti,ab.)) or (HCV adj3 inject*).ti,ab	8407
3	1 and 2	1083
4	limit 3 to (human and yr="2000 -Current" and "review")	93

Skin and soft tissue infections

Medline search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp heroin/ or exp opioid-related disorders/	86581
2	exp Injection site abscess/ or exp Skin Abscess/ or (skin and inject*).ti,ab.	19530
3	1 and 2	256
4	limit 3 to (human and yr="2000 -Current" and "review")	10

Embase search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp opiate/ or exp heroin dependence/	151314
2	exp Injection site abscess/ or exp Skin Abscess/ or (skin and inject*).ti,ab.	40822
3	1 and 2	504
4	limit 3 to (human and yr="2000 -Current" and "review")	25

Quality of life

Medline search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp heroin/ or exp opioid-related disorders/	86581
2	exp Quality of Life/ or "QoL".ti,ab.	162841
3	1 and 2	796
4	limit 3 to (human and yr="2000 -Current" and "review")	153

Embase search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp opiate/ or exp heroin dependence/	151314
2	exp "quality of life"/ or "QoL".ti,ab.	421252
3	1 and 2	4777
4	limit 3 to (human and yr="2000 -Current" and "review")	1163

Contact with criminal justice system

Medline search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp heroin/ or exp opioid-related disorders/	86581

#	Searches	Results
2	exp Offender/ or crim*.ti,ab. or (crim* adj3 behavi*).ti,ab. or (crim* adj3 act*).ti,ab.	30617
3	1 and 2	1041
4	limit 3 to (human and yr="2000 -Current" and "review")	106

Embase search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp opiate/ or exp heroin dependence/	151314
2	exp Offender/ or crim*.ti,ab. or (crim* adj3 behavi*).ti,ab. or (crim* adj3 act*).ti,ab.	50820
3	1 and 2	1666
4	limit 3 to (human and yr="2000 -Current" and "review")	178

Accidental injuries

Medline search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp heroin/ or exp opioid-related disorders/	86581
2	accident\$.ti,ab.	90615
3	1 and 2	500
4	limit 3 to (humans and yr="2000 -Current" and "review")	52

Embase search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp opiate/ or exp heroin dependence/	151314
2	accident\$.ti,ab.	149021
3	1 and 2	1168
4	limit 3 to (human and yr="2000 -Current" and "review")	104

Suicide

Medline search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp heroin/ or exp opioid-related disorders/	86598
2	exp Suicide/ or exp Suicide Attempt/ or suicid*.ti,ab.	75479
3	1 and 2	677
4	limit 3 to (human and yr="2000 -Current" and "review")	51

Embase search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp opiate/ or exp heroin dependence/	151314
2	exp Suicide/ or exp Suicide Attempt/ or suicid*.ti,ab.	107675
3	1 and 2	1566
4	limit 3 to (human and yr="2000 -Current" and "review")	193

Overdose

Medline search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp heroin/ or exp opioid-related disorders/	86581
2	((acute and toxic\$) or overdos\$).ti,ab.	69022
3	1 and 2	2614
4	limit 3 to (humans and yr="2000 -Current" and "review")	340

Embase search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp opiate/ or exp heroin dependence/	151314
2	((acute and toxic\$) or overdos\$).ti,ab.	122589
3	1 and 2	5135
4	limit 3 to (human and yr="2000 -Current" and "review")	528

Neonatal outcomes

Preterm births

Medline search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp heroin/ or exp opioid-related disorders/	86598
2	exp Premature labor/ or exp Prematurity/	22910
3	1 and 2	33
4	limit 3 to (human and yr="2000 -Current" and "review")	3

Embase search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp opiate/ or exp heroin dependence/	151314
2	exp Premature labor/ or exp Prematurity/	136777
3	1 and 2	564
4	limit 3 to (human and yr="2000 -Current" and "review")	130

Low birth weight

Medline search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp heroin/ or exp opioid-related disorders/	86598
2	exp Birth Weight/	38213
3	1 and 2	160
4	limit 3 to (human and yr="2000 -Current" and "review")	5

Embase search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp opiate/ or exp heroin dependence/	151314
2	exp Birth Weight/	109790
3	1 and 2	505
4	limit 3 to (human and yr="2000 -Current" and "review")	56

Head circumference

Medline search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp heroin/ or exp opioid-related disorders/	86598
2	((infant or newborn) adj3 head circumference).ti,ab or (exp Infant/ and head circumference.ti,ab.)	3801
3	1 and 2	60
4	limit 3 to (human and yr="2000 -Current" and "review")	3

Embase search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp opiate/ or exp heroin dependence/	151314
2	exp Head Circumference/	10433
3	1 and 2	89
4	limit 3 to (human and yr="2000 -Current" and "review")	5

Neonatal abstinence syndrome

Medline search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp heroin/ or exp opioid-related disorders/	89598

2	neonatal abstinence syndrome.ti,ab or (exp Infant/ and (withdraw* or abstinence).ti,ab.)	4571
3	1 and 2	621
4	limit 3 to (human and yr="2000 -Current" and "review")	82

Embase search results

#	Searches	Results
1	(opioid or opiate or heroin).ti,ab. or exp opiate/ or exp heroin dependence/	151314
2	exp Neonatal Abstinence Syndrome/	336
3	1 and 2	184
4	limit 3 to (human and yr="2000 -Current" and "review")	33

Description of evidence base on other harms elevated among people who use opioids extra-medically

Injecting risk behaviour

Dependence

HIV incidence

HIV injecting risk

HCV incidence

HCV injecting risk

Skin and soft tissue infections

Quality of life

Criminal activity

Table H1: Self-reported arrests by gender rate ratios reported in Weissman, 1976

Study	Rate ratio	95% confidence interval		% Weight
		Lower	Upper	
Male	4.900	3.900	6.200	85.53
Female	11.400	4.900	26.500	14.47
Pooled rate ratio	5.841	1.358	10.323	100.00




Figure I1: Pooled gender arrests rate ratios as reported in Weissman, 1976.

Contact with criminal justice system

Table H2: Total convictions rate ratios of studies identified via Hayhurst, 2017

Study	Rate ratio	95% confidence interval		% Weight
		Lower	Upper	
Alexander, 1974	0.710	0.400	1.300	20.58
Jarvis, 1989	2.200	1.500	3.100	19.83
Mott, 1974	1.400	1.100	1.700	20.79
Parker, 1987	4.600	3.800	5.600	19.55
Wiepert, 1979	6.200	5.300	7.300	19.25
Pooled rate ratio	2.966	1.426	4.506	100.00




Figure 12: Pooled rate ratios of total convictions of studies identified through Hayhurst, 2017.

Accidental injuries

Suicide

Overdose

Neonatal outcomes

Preterm births

Low birth weight

Head circumference

Neonatal abstinence syndrome

Web appendix I: Details of literature search on mortality among people who use opioids, including fatal overdose

Description of methods

Studies identified in a previously published review will be used to identify studies published from 1980 to 2008.

The Medline, Embase and PsycINFO peer-reviewed literature databases will be searched using the OVID™ interface/platform for relevant articles published from the time-periods 2009 till current. Articles of interest comprise those likely to contain data describing all-cause and cause-specific crude mortality rates (CMR) and/or standardised mortality ratios (SMR) among people who use opioid, cocaine, or amphetamine-type stimulants. Sets of search strings incorporating both keywords and Medical Subject Headings (MeSH terms) reflecting drug type and mortality epidemiology from the previous reviews will be revised and expanded for this updated search. Searches will be limited to human literature. No other restrictions will be applied to the search; citations for papers in languages other than English will be included and read via Google Translate. Citations from these searches will be imported into an Endnote™ library, and duplicate citations removed.

To check for missing peer-review literature, reference lists for relevant systematic reviews identified in the peer-review literature search will be hand searched for additional papers not already identified. A final list of included studies will be distributed to experts to check if any relevant studies are missed.

Each set of search results (title and abstract) will first be screened by one team member in Covidence. All papers marked as excluded will be reviewed by a second person to ensure accuracy in first-pass screening. Each study for full-text screening will be reviewed in full by two people. Conflicts will be resolved through discussion and referral to a third party if needed.

Data will be independently extracted into an Excel worksheet template by one member of the research team and checked by a second member of the team. Bibliographic information will be extracted in addition to study specific information. Data entry will be standardized by use of a

manual, which contains data entry rules. Where data are incomplete, authors will be contacted via email to obtain additional information.

Variables extracted will include study information and sample information (treatment status, HIV status, sex, percentage of sample injecting, treatment engagement). Crude mortality rates (CMRs) and standardized mortality ratios (SMRs) will be extracted as mortality measures. Cause of death information will be extracted for AIDS, overdose, suicide, traumatic (accident, homicide, injury, violence and poisoning), disease-related deaths, and other key categories. Disease-related deaths will be recoded according to the following categories; cardiovascular (endocarditis, myocardial infarction, circulatory system disease and cardiovascular disease), cerebrovascular, respiratory (pneumonia and chronic respiratory disease), liver (cirrhosis, viral hepatitis and liver disease), cancer (neoplasm, tumour and carcinoma), digestive (digestive system disease, nephritis and haemorrhage from duodenal ulcer), nervous system disease and other diseases (tuberculosis, bacterial infection, unspecified 'other disease' or 'natural causes' and when disease categories are not separated).

The quality index used in the previous reviews will be adopted for consistency. The quality index assesses each study on nine individual criteria: case ascertainment, measurement, diagnosis, estimate, numerator and denominator, data catchment, completeness, representativeness and age/sex variables. Each criterion will include a rating scale and the individual scores tallied to provide an overall quality score. Study information necessary for quality assessment will be extracted to the Excel template. The greater the quality score, the higher the methodological quality of the study. Mortality estimates from higher rating studies may be given additional weighting in the calculation of final estimates. Quality estimates may also be used to undertake sensitivity analyses.

Crude mortality rates will be calculated as per 100 person-years. Where person-years are not reported nor made available by the authors, an approximate person-year of follow-up will be calculated, with the assumption that deaths occurred halfway through the follow-up period, so that each case contributes half the person-year follow-up of survivors.

Estimates will be analysed in subgroups according to population type (for example, estimates from cohorts recruited on the basis of having a chronic physical health condition associated with a high mortality rate may be considered separately). Random-effects meta-analyses to determine pooled all-cause and cause-specific CMR and SMR estimates will be performed using STATA. This approach

uses inverse variance weighting to calculate: fixed- and random effects pooled summary estimates; confidence limits; a test for differences between study effects; and an estimate of between-study variance. The random effects model allows for heterogeneity between as well as within studies; expecting high levels of heterogeneity between cohorts, the random-effects model will be used in all meta-analyses, with confirmation through the heterogeneity τ^2 and I-squared statistic. To investigate the source of this heterogeneity in an attempt to reduce it, cohorts may be divided into subsamples (e.g., by sex, age group, treatment status and HIV status) and/or these factors studied as possible risk factors for mortality via meta-regression in Stata.

Search strings for electronic literature searches

Database	Search group	Search terms
Medline*	Opioids	<p>Heroin or opiate or opium or methadone or opioid morphine or buprenorphine or "OST" or "opiate substitution" or "MMT" or "methadone maintenance" or "buprenorphine maintenance" or "opioid replacement" or "opioid agonist" or "opiate agonist" or "opioid maintenance" or "opiate maintenance" or "opiate substitution" or "opiate replacement"</p> <p>exp opium/ or exp narcotics/ or exp heroin dependence/ or exp Heroin/ or exp opioid-related disorders/ or exp opiate alkaloids/ or exp methadone/ or exp analgesics, opioids/ or exp opiate substitution treatment/ or exp buprenorphine/ or exp buprenorphine, naloxone drug combination/</p>
	Mortality	<p>Mortal\$ or fatal\$ or death\$</p> <p>exp DEATH/ or exp "CAUSE OF DEATH"/ or exp DEATH, SUDDEN/ or exp Fatal Outcome/ or exp Mortality/</p>
EMBASE#	Opioids	<p>Heroin or opiate or opium or methadone or opioid or methadone or opioid or morphine or buprenorphine or OST or opioid substitution or MMT or methadone maintenance or buprenorphine maintenance or opioid replacement or opioid agonist or opiate agonist or opioid maintenance or opiate maintenance or opiate substitution or opiate replacement</p> <p>exp Diamorphine/ or exp Opiate/ or exp methadone treatment/ or exp methadone/ or exp heroin dependence/ or exp opiate addiction/ or exp buprenorphine/ or exp buprenorphine plus naloxone/ or exp opiate substitution treatment/</p>
	Mortality	<p>Mortal* or fatal* or death*</p> <p>exp DEATH/ or exp "CAUSE OF DEATH"/ or exp ACCIDENTAL DEATH/ or exp SUDDEN DEATH/ or exp Fatality/ or exp Mortality/</p>
PsychINFO^	Opioids	<p>Heroin or opiate or opium or methadone or opioid or methadone or opioid or morphine or buprenorphine or "OST" or "opioid substitution" or "MMT" or "methadone maintenance" or "buprenorphine maintenance" or "opioid replacement" or "opioid agonist" or "opiate agonist" or "opioid maintenance" or "opiate maintenance" or "opiate substitution" or "opiate replacement"</p> <p>exp Heroin addiction/ or exp Opiates/ or exp methadone/ or exp Heroin/ or exp methadone maintenance/ or exp buprenorphine/</p>
	Mortality	<p>Mortal* or fatal* or death*</p> <p>exp "DEATH AND DYING"/ or exp Mortality/ or exp Mortality Rate</p>

* 'key-words' in lowercase, 'MeSH' terms in **bold**


'key-words' in lowercase, 'EMTREE' terms in **bold**

^ 'key words' in lowercase, explode terms in **bold**

Number of articles identified from opioid mortality search

Search terms			Database		
			EMBASE	Medline	PsycINFO
1.	Opioids	+mortality	8357	3371	1230

PRISMA Flowchart of search strategy



List of included studies

1. Abrahamsson T, Berge J, Ojehagen A, Hakansson A. Benzodiazepine, z-drug and pregabalin prescriptions and mortality among patients in opioid maintenance treatment-A nation-wide register-based open cohort study. *Drug and Alcohol Dependence* 2017; **174**: 58-64.
2. Antolini G, Pirani M, Morandi G, Sorio C. Gender difference and mortality in a cohort of heroin users in the Provinces of Modena and Ferrara, 1975-1999. *Epidemiologia e Prevenzione* 2006; **30**(2): 91-9.
3. Apelt SM, Scherbaum N, Golz J, Backmund M, Soyka M. Safety, effectiveness and tolerance of buprenorphine-naloxone in the treatment of opioid dependence: Results from a nationwide non-interventional study in routine care. *Pharmacopsychiatry* 2013; **46**(3): 94-107.
4. Arendt M, Munk-Jorgensen P, Sher L, Jensen SO. Mortality among individuals with cannabis, cocaine, amphetamine, MDMA, and opioid use disorders: A nationwide follow-up study of Danish substance users in treatment. *Drug and Alcohol Dependence* 2011; **114**(2-3): 134-9.
5. Azim T, Chowdhury EI, Reza M, et al. Prevalence of infections, HIV risk behaviors and factors associated with HIV infection among male injecting drug users attending a needle/syringe exchange program in Dhaka, Bangladesh. *Subst Use Misuse* 2008; **43**(14): 2124-44.
6. Bakker A, Streel E. Benzodiazepine maintenance in opiate substitution treatment: Good or bad? A retrospective primary care case-note review. *Journal of Psychopharmacology* 2017; **31**(1): 62-6.
7. Bargagli AM, Hickman M, Davoli M, et al. Drug-related mortality and its impact on adult mortality in eight European countries. *European Journal of Public Health* 2006; **16**(2): 198-202.
8. Bargagli AM, Sperati A, Davoli M, Forastiere F, Perucci CA. Mortality among problem drug users in Rome: an 18-year follow-up study, 1980-97. *Addiction* 2001; **96**(10): 1455-63.
9. Barrio G, Molist G, de la Fuente L, et al. Mortality in a cohort of young primary cocaine users: Controlling the effect of the riskiest drug-use behaviors. *Addictive Behaviors* 2013; **38**(3): 1601-4.
10. Bart G, Wyman Z, Wang Q, Hodges JS, Karim R, Bart BA. Methadone and the QTc Interval: Paucity of Clinically Significant Factors in a Retrospective Cohort. *Journal of Addiction Medicine* 2017; **11**(6): 489-93.
11. Bartu A. Bartu et al. (2004): A correction. *Addiction* 2006; **101**(10): 1524.
12. Bartu A, Freeman NC, Gawthorne GS, Codde JP, Holman CDJ. Mortality in a cohort of opiate and amphetamine users in Perth, Western Australia. *Addiction* 2004; **99**(1): 53-60.
13. Bauer SM, Loipl R, Jagsch R, et al. Mortality in opioid-maintained patients after release from an addiction clinic. *European Addiction Research* 2008; **14**(2): 82-91.
14. Bogdanowicz KM, Stewart R, Broadbent M, et al. Double trouble: Psychiatric comorbidity and opioid addiction-All-cause and cause-specific mortality. *Drug and Alcohol Dependence* 2015; **148**: 85-92.
15. Bogdanowicz KM, Stewart R, Chang C-K, et al. Identifying mortality risks in patients with opioid use disorder using brief screening assessment: Secondary mental health clinical records analysis. *Drug and Alcohol Dependence* 2016; **164**: 82-8.
16. Brugal MT, Molist G, Sarasa-Renedo A, et al. Assessing gender disparities in excess mortality of heroin or cocaine users compared to the general population. *International Journal of Drug Policy* 2016; **38**: 36-42.
17. Buster MCA, van Brussel GHA, van den Brink W. An increase in overdose mortality during the first 2 weeks after entering or re-entering methadone treatment in Amsterdam. *Addiction* 2002; **97**(8): 993-1001.
18. Callaghan RC, Cunningham JK, Verdichevski M, Sykes J, Jaffer SR, Kish SJ. All-cause mortality among individuals with disorders related to the use of methamphetamine: A comparative cohort study. *Drug and Alcohol Dependence* 2012; **125**(3): 290-4.
19. Callaghan RC, Gatley JM, Veldhuizen S, Lev-Ran S, Mann R, Asbridge M. Alcohol- or drug-use disorders and motor vehicle accident mortality: A retrospective cohort study. *Accident Analysis and Prevention* 2013; **53**: 149-55.
20. Chang K-C, Lu T-H, Lee K-Y, Hwang J-S, Cheng C-M, Wang J-D. Estimation of life expectancy and the expected years of life lost among heroin users in the era of opioid substitution treatment (OST) in Taiwan. *Drug and Alcohol Dependence* 2015; **153**: 152-8.
21. Chang K-C, Wang J-D, Saxon A, Matthews AG, Woody G, Hser Y-I. Causes of death and expected years of life lost among treated opioid-dependent individuals in the United States and Taiwan. *International Journal of Drug Policy* 2017; **43**: 1-6.

22. Chen CY, Wu PN, Su LW, Chou YJ, Lin KM. Three-year mortality and predictors after release: a longitudinal study of the first-time drug offenders in Taiwan. *Addiction* 2010; **105**(5): 920-7.
23. Clausen T, Anchersen K, Waal H. Mortality prior to, during and after opioid maintenance treatment (OMT): A national prospective cross-registry study. *Drug and Alcohol Dependence* 2008; **94**(1-3): 151-7.
24. Clausen T, Waal H, Thoresen M, Gossop M. Mortality among opiate users: Opioid maintenance therapy, age and causes of death. *Addiction* 2009; **104**(8): 1356-62.
25. Cornish R, Macleod J, Strang J, Vickerman P, Hickman M. Risk of death during and after opiate substitution treatment in primary care: Prospective observational study in UK General Practice Research Database. *BMJ: British Medical Journal* 2010; **341**(7779): 1-8.
26. Cottler LB, Hu H, Smallwood BA, Anthony JC, Wu LT, Eaton WW. Nonmedical Opioid Pain Relievers and All-Cause Mortality: A 27-Year Follow-Up From the Epidemiologic Catchment Area Study. *American journal of public health* 2016; **106**(3): 509-16.
27. Dasgupta N, Funk MJ, Proescholdbell S, Hirsch A, Ribisl KM, Marshall S. Cohort study of the impact of high-dose opioid analgesics on overdose mortality. *Pain Medicine (United States)* 2016; **17**(1): 85-98.
28. Davstad I, Allebeck P, Leifman A, Stenbacka M, Romelsjo A. Self-reported drug use and mortality among a nationwide sample of Swedish conscripts - A 35-year follow-up. *Drug and Alcohol Dependence* 2011; **118**(2-3): 383-90.
29. de la Fuente L, Molist G, Espelt A, et al. Mortality risk factors and excess mortality in a cohort of cocaine users admitted to drug treatment in Spain. *Journal of Substance Abuse Treatment* 2014; **46**(2): 219-26.
30. Degenhardt L, Larney S, Kimber J, et al. The impact of opioid substitution therapy on mortality post-release from prison: Retrospective data linkage study. *Addiction* 2014; **109**(8): 1306-17.
31. Degenhardt L, Larney S, Randall D, Burns L, Hall W. Causes of death in a cohort treated for opioid dependence between 1985 and 2005. *Addiction* 2014; **109**(1): 90-9.
32. Degenhardt L, Randall D, Hall W, Law M, Butler T, Burns L. Mortality among clients of a state-wide opioid pharmacotherapy program over 20 years: Risk factors and lives saved. *Drug and Alcohol Dependence* 2009; **105**(1-2): 9-15.
33. Delorme J, Chenaf C, Kabore J-L, et al. Incidence of high dosage buprenorphine and methadone shopping behavior in a retrospective cohort of opioid-maintained patients in France. *Drug and Alcohol Dependence* 2016; **162**: 99-106.
34. Dunn KM, Saunders KW, Rutter CM, et al. Opioid prescriptions for chronic pain and overdose: A cohort study. *Annals of Internal Medicine* 2010; **152**(2): 85-92.
35. Ekholm O, Kurita GP, Hojsted J, Juel K, Sjogren P. Chronic pain, opioid prescriptions, and mortality in Denmark: A population-based cohort study. *Pain* 2014; **155**(12): 2486-90.
36. Espelt A, Barrio G, Alamo-Junquera D, et al. Lethality of opioid overdose in a community cohort of young heroin users. *European Addiction Research* 2015; **21**(6): 300-6.
37. Esteban J, Gimeno C, Barril J, Aragonés A, Climent JM, de la Cruz Pellin M. Survival study of opioid addicts in relation to its adherence to methadone maintenance treatment. *Drug and Alcohol Dependence* 2003; **70**(2): 193-200.
38. Evans E, Li L, Min J, et al. Mortality among individuals accessing pharmacological treatment for opioid dependence in California, 2006-10. *Addiction* 2015; **110**(6): 996-1005.
39. Evans E, Li L, Min JE, et al. Gender differences in mortality among treated opioid-dependent patients. *Drug and Alcohol Dependence* 2015; **156**: e64.
40. Evans JL, Tsui JI, Hahn JA, Davidson PJ, Lum PJ, Page K. Mortality among young injection drug users in San Francisco: A 10-year follow-up of the UFO study. *American Journal of Epidemiology* 2012; **175**(4): 302-8.
41. Fellows-Smith J. Opioid-dependent error processing. *Journal of Opioid Management* 2011; **7**(6): 443-9.
42. Ferri M, Bargagli AM, Faggiano F, et al. Mortality of drug users attending public treatment centres in Italy 1998-2001: A cohort study. *Epidemiologia e Prevenzione* 2007; **31**(5): 276-82.
43. Foster D, Udayachalerm S, Wang J, Murray M. Characterization of opioid use and adverse outcomes using longitudinal data from a statewide health information exchange. *Pharmacotherapy* 2017; **37** (12): e189.
44. Fridell M, Hesse M. Psychiatric severity and mortality in substance abusers: A 15-year follow-up of drug users. *Addictive Behaviors* 2006; **31**(4): 559-65.

45. Fugelstad A, Agren G, Romelsjö A. Changes in Mortality, Arrests, and Hospitalizations in Nonvoluntary Treated Heroin Addicts in Relation to Methadone Treatment. *Substance Use & Misuse* 1998; **33**(14): 2803-17.
46. Fugelstad A, Annell A, Agren G. Long-term mortality and causes of death among hospitalized Swedish drug users. *Scandinavian journal of public health* 2014; **42**(4): 364-9.
47. Galli M, Musicco M. Mortality of intravenous drug users living in Milan, Italy: Role of HIV-1 infection. *AIDS* 1994; **1994**(8): 1457-63.
48. Gao L, Dimitropoulou P, Robertson JR, McTaggart S, Bennie M, Bird SM. Risk-factors for methadone-specific deaths in Scotland's methadone-prescription clients between 2009 and 2013. *Drug and Alcohol Dependence* 2016; **167**: 214-23.
49. Garg RK, Fulton-Kehoe D, Franklin GM. Patterns of opioid use and risk of opioid overdose death among Medicaid patients. *Medical Care* 2017; **55**(7): 661-8.
50. Ghodse H, Oyefeso A, Kilpatrick B. Mortality of drug addicts in the United Kingdom 1967-1993. *International Journal of Epidemiology* 1998; **27**(3): 473-8.
51. Gibson A, Randall D, Degenhardt L. The increasing mortality burden of liver disease among opioid-dependent people: Cohort study. *Addiction* 2011; **106**(12): 2186-92.
52. Gjersing L, Bretteville-Jensen AL. Gender differences in mortality and risk factors in a 13-year cohort study of street-recruited injecting drug users. *BMC public health* 2014; **14**: 440.
53. Goldstein A, Herrera J. Heroin addicts and methadone treatment in Albuquerque: A 22-year follow up. *Drug and Alcohol Dependence* 1995; **40**(2): 139-50.
54. Haarr D, Nessa J. Treatment of opiate-dependent patients in a general practice. *Tidsskr for den Norske Laegeforening: Tidsskrift for Praktisk Medicin, ny Raekke* 2007; **127**(13): 1770-2.
55. Hayashi K, Dong H, Marshall BDL, et al. Sex-Based Differences in Rates, Causes, and Predictors of Death among Injection Drug Users in Vancouver, Canada. *American Journal of Epidemiology* 2016; **183**(6): 544-52.
56. Hser Y-I, Mooney LJ, Saxon AJ, et al. High mortality among patients with opioid use disorder in a large healthcare system. *Journal of Addiction Medicine* 2017; **11**(4): 315-9.
57. Huang CL-C, Master CWL. Factors associated with mortality among heroin users after seeking treatment with methadone: A population-based cohort study in Taiwan. *Journal of Substance Abuse Treatment* 2013; **44**(3): 295-300.
58. Huang Y-F, Kuo H-S, Lew-Ting C-Y, et al. Mortality among a cohort of drug users after their release from prison: An evaluation of the effectiveness of a harm reduction program in Taiwan. *Addiction* 2011; **106**(8): 1437-45.
59. Jafari S, Rahimi-Movaghar A, Craib KJP, Baharlou S, Mathias R. A follow-up study of drug users in Southern Iran. *Addiction Research & Theory* 2010; **18**(1): 59-70.
60. Jerkeman A, Hakansson A, Rylance R, Wagner P, Blome MA, Bjorkman P. Death from liver disease in a cohort of injecting opioid users in a Swedish city in relation to registration for opioid substitution therapy. *Drug and Alcohol Review* 2017; **36**(3): 424-31.
61. Jimenez-Trevino L, Saiz PA, Garcia-Portilla MP, et al. A 25-year follow-up of patients admitted to methadone treatment for the first time: Mortality and gender differences. *Addictive Behaviors* 2011; **36**(12): 1184-90.
62. Jones AA, Vila-Rodriguez F, Leonova O, et al. Mortality from treatable illnesses in marginally housed adults: a prospective cohort study. *BMJ Open* 2015; **5**(8): e008876.
63. Kelty E, Dobbins T, Hulse G. Incidence of cancer and cancer related mortality in opiate dependent patients treated with methadone, buprenorphine or implant naltrexone as compared with non-opiate using controls. *Heroin Addiction and Related Clinical Problems* 2017; **19**(3): 65-72.
64. Kelty E, Hulse G. Examination of mortality rates in a retrospective cohort of patients treated with oral or implant naltrexone for problematic opiate use. *Addiction* 2012; **107**(10): 1817-24.
65. Kelty E, Hulse G. Fatal and non-fatal opioid overdose in opioid dependent patients treated with methadone, buprenorphine or implant naltrexone. *International Journal of Drug Policy* 2017; **46**: 54-60.
66. Khademi H, Malekzadeh R, Pourshams A, et al. Opium use and mortality in Golestan Cohort Study: prospective cohort study of 50,000 adults in Iran. *BMJ (Clinical research ed)* 2012; **344**(e2502): 1-12.

67. Krebs EE, Becker WC, Zerzan J, Bair MJ, McCoy K, Hui S. Comparative mortality among Department of Veterans Affairs patients prescribed methadone or long-acting morphine for chronic pain. *Pain* 2011; **152**(8): 1789-95.
68. Langendam MW, van Brussel GHA, Coutinho RA, van Ameijden EJC. The impact of harm-reduction-based methadone treatment on mortality among heroin users. *American Journal of Public Health* 2001; **91**(5): 774-80.
69. Larney S, Bohnert AS, Ganoczy D, et al. Mortality among older adults with opioid use disorders in the Veteran's Health Administration, 2000-2011. *Drug and Alcohol Dependence* 2015; **147**: 32-7.
70. Larney S, Randall D, Gibson A, Degenhardt L. The contributions of viral hepatitis and alcohol to liver-related deaths in opioid-dependent people. *Drug and Alcohol Dependence* 2013; **131**(3): 252-7.
71. Laroche MR, Bernson D, Land T, et al. Medication for Opioid Use Disorder After Nonfatal Opioid Overdose and Association With Mortality: A Cohort Study. *Ann Intern Med* 2018.
72. Ledberg A. Mortality related to methadone maintenance treatment in Stockholm, Sweden, during 2006-2013. *Journal of Substance Abuse Treatment* 2017; **74**: 35-41.
73. Lee CTC, Chen VCH, Tan HKL, et al. Suicide and other-cause mortality among heroin users in Taiwan: A prospective study. *Addictive Behaviors* 2013; **38**(10): 2619-23.
74. Lejckova P, Mravcik V. Mortality of hospitalized drug users in the Czech Republic. *Journal of Drug Issues* 2007; **37**(1): 103-18.
75. Liao D-L, Chen P-C, Chen C-H, et al. Higher methadone doses are associated with lower mortality in patients of opioid dependence in Taiwan. *Journal of Psychiatric Research* 2013; **47**(10): 1530-4.
76. Lindblad R, Hu L, Oden N, Wakim P, Rosa C, VanVeldhuisen P. Mortality Rates Among Substance Use Disorder Participants in Clinical Trials: Pooled Analysis of Twenty-Two Clinical Trials Within the National Drug Abuse Treatment Clinical Trials Network. *Journal of Substance Abuse Treatment* 2016; **70**: 73-80.
77. Liu E, Rou K, McGoogan JM, et al. Factors associated with mortality of HIV-positive clients receiving methadone maintenance treatment in China. *Journal of Infectious Diseases* 2013; **208**(3): 442-53.
78. Lloyd B, Barratt MJ, Ferris J, Best D, Lubman DI. Factors influencing mortality among alcohol and drug treatment clients in Victoria, Australia: The role of demographic and substance use characteristics. *Australian and New Zealand Journal of Psychiatry* 2013; **47**(9): 859-67.
79. Lopez D, Martineau H, Palle C. Mortality of individuals arrested for heroin, cocaine or crack use. Paris, France: Observatoire Francais des Drogues et des Toxicomanies, 2004.
80. Lopez-Quintero C, Roth KB, Eaton WW, et al. Mortality among heroin users and users of other internationally regulated drugs: A 27-year follow-up of users in the Epidemiologic Catchment Area Program household samples. *Drug and Alcohol Dependence* 2015; **156**: 104-11.
81. Lovrecic B, Semerl JS, Tavcar R, Maremmani I. Sociodemographic and clinical differences among deceased and surviving cohort members of opioid maintenance therapy. *Heroin Addiction and Related Clinical Problems* 2011; **13**(3): 39-48.
82. Marsden J, Stillwell G, Jones H, et al. Does exposure to opioid substitution treatment in prison reduce the risk of death after release? A national prospective observational study in England. *Addiction* 2017; **112**(8): 1408-18.
83. Marzo J-N, Rotily M, Meroueh F, et al. Maintenance therapy and 3-year outcome of opioid-dependent prisoners: A prospective study in France (2003-06). *Addiction* 2009; **104**(7): 1233-40.
84. Merrill ELC, Bird SM, Hutchinson SJ. Mortality of those who attended drug services in Scotland 1996-2006: Record-linkage study. *International Journal of Drug Policy* 2012; **23**(1): 24-32.
85. Metzger DS, Donnell D, Celentano DD, et al. Expanding substance use treatment options for HIV prevention with buprenorphine-naloxone: HIV Prevention Trials Network 058. *JAIDS Journal of Acquired Immune Deficiency Syndromes* 2015; **68**(5): 554-61.
86. Molist G, Brugal MT, Barrio G, et al. Effect of ageing and time since first heroin and cocaine use on mortality from external and natural causes in a Spanish cohort of drug users. *International Journal of Drug Policy* 2018; **53**: 8-16.
87. Moskalewicz J, Sieroslawski J. Mortality of narcotic addicts using injections. *Przegląd Epidemiologiczny* 1996; **50**(3): 323-32.

88. Muga R, Rivas I, Faure E, et al. Sex-specific disease outcomes of HIV-positive and HIV-negative drug users admitted to an opioid substitution therapy program in Spain: A cohort study. *BMC Infectious Diseases* 2014; **14** (504).
89. Muhuri PK, Gfroerer JC. Mortality associated with illegal drug use among adults in the United States. *The American Journal of Drug and Alcohol Abuse* 2011; **37**(3): 155-64.
90. Naderi-Heiden A, Gleiss A, Backer C, et al. Mortality and employment after in-patient opiate detoxification. *European Psychiatry* 2012; **27**(4): 294-300.
91. Nagot N, Hoang TG, Vallo R, et al. Mortality among people who inject drugs in Haiphong, Vietnam: High burden of infectious causes. 2018.
92. Nambiar D, Agius PA, Stooze M, Hickman M, Dietze P. Mortality in the Melbourne injecting drug user cohort study (MIX). *Harm Reduction Journal* 2015; **12**(55): 1-5.
93. O'Connor G, McGinty T, Yeung SJ, et al. Cross-sectional study of the characteristics, healthcare usage, morbidity and mortality of injecting drug users attending an inner city emergency department. *Emergency Medicine Journal* 2014; **31**(8): 625-9.
94. Odegard E, Amundsen EJ, Kielland KB. Fatal overdoses and deaths by other causes in a cohort of Norwegian drug abusers--a competing risk approach. *Drug and Alcohol Dependence* 2007; **89**(2-3): 176-82.
95. Oppenheimer E, Tobutt C, Taylor C, Andrew T. Death and survival in a cohort of heroin addicts from London clinics: A 22-year follow-up study. *Addiction* 1994; **89**(10): 1299-308.
96. Orti RM, Domingo-Salvany A, Munoz A, MacFarlane D, Suelves JM, Anto JM. Mortality trends in a cohort of opiate addicts, Catalonia, Spain. *International Journal of Epidemiology* 1996; **25**(3): 545-53.
97. Pan C-H, Jhong J-R, Tsai S-Y, Lin S-K, Chen C-C, Kuo C-J. Excessive suicide mortality and risk factors for suicide among patients with heroin dependence. *Drug and Alcohol Dependence* 2014; **145**: 224-30.
98. Parmar MK, Strang J, Choo L, Meade AM, Bird SM. Randomized controlled pilot trial of naloxone-on-release to prevent post-prison opioid overdose deaths. *Addiction* 2017; **112**(3): 502-15.
99. Pavarin RM, Fioritti A, Sanchini S. Mortality trends among heroin users treated between 1975 and 2013 in Northern Italy: Results of a longitudinal study. *Journal of Substance Abuse Treatment* 2017; **77**: 166-73.
100. Peles E, Schreiber S, Adelson M. 15-year survival and retention of patients in a general hospital-affiliated methadone maintenance treatment (MMT) center in Israel. *Drug and Alcohol Dependence* 2010; **107**(2-3): 141-8.
101. Peles E, Schreiber S, Adelson M. Opiate-dependent patients on a waiting list for methadone maintenance treatment are at high risk for mortality until treatment entry. *Journal of Addiction Medicine* 2013; **7**(3): 177-82.
102. Pierce M, Bird SM, Hickman M, Millar T. National record linkage study of mortality for a large cohort of opioid users ascertained by drug treatment or criminal justice sources in England, 2005-2009. *Drug and Alcohol Dependence* 2015; **146**: 17-23.
103. Price RK, Risk NK, Murray KS, Virgo KS, Spitznagel EL. Twenty-five year mortality of US serviceman deployed in Vietnam: Predictive utility of early drug use. *Drug and Alcohol Dependence* 2001; **64**(3): 309-18.
104. Quan VM, Minh NL, Ha TV, et al. Mortality and HIV transmission among male Vietnamese injection drug users. *Addiction* 2011; **106**(3): 583-9.
105. Quan VM, Vongchak T, Jittiwutikarn J, et al. Predictors of mortality among injecting and non-injecting HIV-negative drug users in northern Thailand. *Addiction* 2007; **102**(3): 441-6.
106. Ray WA, Chung CP, Murray KT, Hall K, Michael Stein C. Prescription of long-acting opioids and mortality in patients with chronic noncancer pain. *JAMA - Journal of the American Medical Association* 2016; **315**(22): 2415-23.
107. Reece AS. Favorable mortality profile of naltrexone implants for opiate addiction. *Journal of Addictive Diseases* 2010; **29**(1): 30-50.
108. Risser D, Honigschnabl S, Stichenwirth M, et al. Mortality of opiate users in Vienna, Austria. *Drug and Alcohol Dependence* 2001; **64**(3): 251-6.
109. Rossow I. Suicide among drug addicts in Norway. *Addiction* 1994; **89**(12): 1667-73.
110. Sanchez-Carbonell X, Seus L. Ten-year survival analysis of a cohort of heroin addicts in Catalonia: The EMETYST project *Addiction* 2000; **95**(6): 941-8.

111. Sanvisens A, Vallecillo G, Bolao F, et al. Temporal trends in the survival of drug and alcohol abusers according to the primary drug of admission to treatment in Spain. *Drug and Alcohol Dependence* 2014; **136**: 115-20.
112. Scherbaum N, Specka M, Hauptmann G, Gastpar M. Does maintenance treatment reduce the mortality rate of opioid addicts? *Fortschritte der Neurologie-Psychiatrie* 2002; **70**(9): 455-61.
113. Smyth BP, Fagan J, Kernan K. Outcome of heroin-dependent adolescents presenting for opiate substitution treatment. *Journal of Substance Abuse Treatment* 2012; **42**(1): 35-44.
114. Solomon DH, Rassen JA, Glynn RJ, et al. The comparative safety of opioids for nonmalignant pain in older adults. *Archives of Internal Medicine* 2010; **170**(22): 1979-86.
115. Solomon DH, Rassen JA, Glynn RJ, Lee J, Levin R, Schneeweiss S. The comparative safety of analgesics in older adults with arthritis. *Archives of Internal Medicine* 2010; **170**(22): 1968-78.
116. Soyka M, Apelt SM, Lieb M, Wittchen HU. One-year mortality rates of patients receiving methadone and buprenorphine maintenance therapy: a nationally representative cohort study in 2694 patients. *Journal of Clinical Psychopharmacology* 2006; **26**(6): 657-60.
117. Soyka M, Trader A, Klotsche J, et al. Six-year mortality rates of patients in methadone and buprenorphine maintenance therapy: Results from a nationally representative cohort study. *Journal of Clinical Psychopharmacology* 2011; **31**(5): 678-80.
118. Stenbacka M, Leifman A, Romelsjo A. Mortality among opiate abusers in Stockholm: A longitudinal study. *Heroin Addiction and Related Clinical Problems* 2007; **9**(3): 41-50.
119. Stooze MA, Dietze PM, Jolley D. Overdose deaths following previous non-fatal heroin overdose: Record linkage of ambulance attendance and death registry data. *Drug and alcohol review* 2009; **28**(4): 347-52.
120. Tait RJ, Ngo HT, Hulse GK. Mortality in heroin users 3 years after naltrexone implant or methadone maintenance treatment. *Journal of Substance Abuse Treatment* 2008; **35**(2): 116-24.
121. Uosukainen H, Kauhanen J, Bell J, et al. Mortality among clients seeking treatment for buprenorphine abuse in Finland. *Drug and Alcohol Dependence* 2013; **133**(2): 391-7.
122. Veldhuizen S, Callaghan RC. Cause-specific mortality among people previously hospitalized with opioid-related conditions: A retrospective cohort study. *Annals of Epidemiology* 2014; **24**(8): 620-4.
123. Vlahov D, Wang C, Ompad D, et al. Mortality risk among recent-onset injection drug users in five U.S. cities. *Substance Use & Misuse* 2008; **43**(3-4): 413-28.
124. Wahren CA, Brandt L, Allebeck P. Has mortality in drug addicts increased? A comparison between two hospitalized cohorts in Stockholm. *International Journal of Epidemiology* 1997; **26**(6): 1219-26.
125. Wang C, Vlahov D, Galai N, et al. The effect of HIV infection on overdose mortality. *AIDS* 2005; **19**(9): 935-42.
126. Zhang L, Ruan YH, Jiang ZQ, et al. A 1-year prospective cohort study on mortality of injecting drug users. *Zhonghua liu xing bing xue za zhi = Zhonghua liuxingbingxue zazhi* 2005; **26**(3): 190-3.

Web appendix J: Details of literature search on reviews of interventions to reduce adverse health outcomes

Description of Method

We used a multi-stage hierarchical searching process to identify sources reporting on the impact of various interventions on health outcomes associated with amphetamine and cocaine use amongst samples who report use of these substances.

The intent was to search for and identify the highest quality study type (see Table 1) for each intervention and the associated health outcome.

Table 1. Classification system used in assessment of study methodologies

Level	Study type
1	Systematic review of all relevant randomised controlled trials
2	Properly designed randomised controlled trial
3.1	Well-designed pseudorandomised controlled trials [alternate allocation or some other method]
3.2	Comparative studies with concurrent controls and allocation not randomised [cohort studies], case-control studies, or interrupted time series with a control group
4	Case series, pre–test, and post–test, cross-sectional

There were four stages to searching, with stages 2-4 only undertaken if the highest level study type (i.e., level 1: systematic review of randomised controlled trials) was not identified in the previous stage. The hierarchy for selection of studies is detailed in Table 1. It should be noted that where narrative reviews were identified in stage 1 and 2, reference lists were searched for high-quality studies. Exclude study type 4.

Table 2. Hierarchy of searching for studies

Stage	Details
1	Cochrane Library of Systematic Reviews and EBM reviews search for Level 1 studies
2	Peer-reviewed literature database (Medline, Embase, PsycINFO) search for Level 1 studies
3	Peer-reviewed literature database (Medline, Embase, PsycINFO) search for Level 2-4 studies

Search Strategy**Cochrane Library of Systematic Reviews and EBM reviews (Full Text-Cochrane DSR, ACP Journal Club and DARE) search for Level 1 studies**

The following searches were limited to those references published since 2000 and only Cochrane and CRD reviews.

Cochrane

	Search Group	Number of citations retrieved
1	opioid OR opiate OR heroin	16845
2	Limit 1 to "Cochrane reviews" and "2000-2018"	205

EBM

	Search Group	Number of citations retrieved
1	opioid.mp. [mp=title, short title, abstract, full text, keywords, caption text]	464
2	opiate.mp. [mp=title, short title, abstract, full text, keywords, caption text]	180
3	heroin.mp. [mp=title, short title, abstract, full text, keywords, caption text]	93
4	1 or 2 or 3	554
5	limit 4 to (full systematic reviews and last 18 years)	421

Peer-reviewed literature database searches for Level 1 studies

Medline search results

#	Terms	Results
1	heroin*.ti,ab.	11849
2	opiate*.ti,ab.	22313
3	opium*.ti,ab.	1687
4	opioid*.ti,ab.	64936
5	morphine*.ti,ab.	43307
6	exp opium/	1944
7	exp narcotics/	111411
8	exp heroin dependence/	8659
9	exp Heroin/	5251
10	exp opioid-related disorders/	22435
11	exp opiate alkaloids/	81411
12	exp analgesics, opioid/	103531
13	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12	177000
14	exp intravenous drug abuse/	13925
15	injecting risk.ti,ab.	129
16	"IDU\$1".tw.	5019
17	"IVDU\$1".tw.	718
18	"PWID\$1".tw.	718
19	"injecting drug".tw.	3694
20	"intravenous drug".tw.	7165
21	"injecting substance".tw.	9
22	"intravenous substance".tw.	56
23	exp substance abuse, intravenous/	13925
24	14 or 15 or 16 or 17 or 18 or 19 or 20 or 21 or 22 or 23	21532

#	Terms	Results
25	13 or 24	194862
26	exp intervention study/	790089
27	exp early intervention/	2495
28	drug dependence/di, dt, pc, rh, th [Diagnosis, Drug Therapy, Prevention, Rehabilitation, Therapy]	34941
29	exp substance abuse treatment centers/	5000
30	"drug dependence treatment".ti,ab.	100
31	exp harm reduction/	2347
32	Addiction/dt [Drug Therapy]	0
33	26 or 27 or 28 or 29 or 30 or 31 or 32	829093
34	(brief adj3 intervention).ti,ab.	3584
35	(safe adj3 inject*).ti,ab.	877
36	(needle adj2 syringe adj5 program).ti,ab.	66
37	exp condom/	9356
38	sexually transmitted disease/di, dt, pc, rh, th [Diagnosis, Drug Therapy, Prevention, Rehabilitation, Therapy]	12513
39	hepatitis C/di, dt, pc, rh, th [Diagnosis, Drug Therapy, Prevention, Rehabilitation, Therapy]	16277
40	Human immunodeficiency virus/pc [Prevention]	0
41	exp pre-exposure prophylaxis/	961
42	(prep adj5 HIV).ti,ab.	532
43	(prep adj5 sexual*).ti,ab.	72
44	(pre-exposure prophylaxis adj5 sexual*).mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]	109
45	(drug adj3 consum* adj3 (room or facil*)).ti,ab.	25
46	(supervis* adj3 inject*).ti,ab.	226

#	Terms	Results
47	atypical antipsychotic agent/dt [Drug Therapy]	0
48	Suicide/pc [Prevention]	8402
49	(suicide adj3 prevent*).ti,ab.	4249
50	exp behavior therapy/	65237
51	exp motivational interviewing/	1222
52	exp cognitive therapy/	23038
53	"cognitive behavio?r therapy".ti,ab.	2635
54	exp self-help groups/	9629
55	((peer or mutual) adj3 support*).ti,ab.	3835
56	exp family therapy/	8414
57	multisystemic therapy.ti,ab.	156
58	exp therapeutic community/	2116
59	agonist pharmacotherap*.ti,ab.	22
60	(drug adj3 (compulsory or detention)).ti,ab.	95
61	"NSP\$1".tw.	4021
62	"needle syringe program\$".tw.	47
63	"NSEP\$1".tw.	56
64	"needle syringe exchange program\$".tw.	23
65	"needle exchange\$1".tw.	871
66	"syringe exchange\$1".tw.	569
67	exp Needle-Exchange Programs/	1551
68	exp Harm Reduction/	2347
69	"HAART".tw.	10467
70	"Anti?Retroviral Treatment".tw.	6218
71	"Anti?Retroviral Therapy".tw.	31258
72	"highly active anti?retroviral therapy".tw.	9978

#	Terms	Results
73	"highly active anti?retroviral treatment".tw.	491
74	"HIV treatment".tw.	3291
75	"anti-HIV Agents".tw.	757
76	"AIDS treatment".tw.	939
77	exp anti-hiv agents/	61737
78	exp hiv fusion inhibitors/	1136
79	exp hiv integrase inhibitors/	2139
80	exp hiv protease inhibitors/	12338
81	exp Antiretroviral Therapy, Highly Active/	19564
82	"OST".tw.	1066
83	"opioid substitution treatment".tw.	198
84	"methadone".tw.	11139
85	"MMT".tw.	1992
86	"methadone maintenance".tw.	3526
87	"buprenorphine".tw.	4856
88	"BMT".tw.	9731
89	"Buprenorphine maintenance".tw.	224
90	"opioid replacement".tw.	96
91	exp Buprenorphine/	4529
92	exp Methadone/	11561
93	exp Opiate Substitution Treatment/	2052
94	exp Buprenorphine, Naloxone Drug Combination/	190
95	"naltrexone".tw.	5769
96	"LAAM".tw.	293
97	"levomethadyl acetate".tw.	20
98	exp Naltrexone/	7339

#	Terms	Results
99	exp Methadyl Acetate/	406
100	"contingency management".tw.	754
101	Vaccine/dt [Drug Therapy]	0
102	"detoxification".ti,ab.	23495
103	"drug detoxification".ti,ab.	296
104	exp social support/	62877
105	exp rehabilitation centers/	13595
106	exp "acceptance and commitment therapy"/	216
107	34 or 35 or 36 or 37 or 38 or 39 or 40 or 41 or 42 or 43 or 44 or 45 or 46 or 47 or 48 or 49 or 50 or 51 or 52 or 53 or 54 or 55 or 56 or 57 or 58 or 59 or 60 or 61 or 62 or 63 or 64 or 65 or 66 or 67 or 68 or 69 or 70 or 71 or 72 or 73 or 74 or 75 or 76 or 77 or 78 or 79 or 80 or 81 or 82 or 83 or 84 or 85 or 86 or 87 or 88 or 89 or 90 or 91 or 92 or 93 or 94 or 95 or 96 or 97 or 98 or 99 or 100 or 101 or 102 or 103 or 104 or 105 or 106	359004
108	33 or 107	1138477
109	exp Evaluation Studies/	233034
110	exp program evaluation/	67455
111	exp evidence based medicine/	67673
112	exp "Outcome Assessment (Health Care)"/	921018
113	exp patient outcome assessment/	4985
114	exp randomized controlled trial/	455867
115	random* controlled trial.ti,ab.	61699
116	exp Clinical Trials as Topic/	311580
117	exp controlled clinical trial/	543249
118	multicenter study/	230085
119	random*.ti,ab.	825940
120	(pretest or pre test).ti,ab.	13379
121	(posttest or post test).ti,ab.	13989

#	Terms	Results
122	before after.ti,ab.	3518
123	qua?irandomi*.ti,ab.	56
124	nonrandomi*.ti,ab.	9746
125	exp cohort analysis/	1720362
126	exp longitudinal study/	113428
127	exp Meta-Analysis as Topic/	16350
128	systematic review/	0
129	exp Clinical Trial/	790089
130	109 or 110 or 111 or 112 or 113 or 114 or 115 or 116 or 117 or 118 or 119 or 120 or 121 or 122 or 123 or 124 or 125 or 126 or 127 or 128 or 129	3593803
131	25 and 108 and 130	28470
132	limit 131 to (humans and yr="2000 -Current" and "review")	796

Searches conducted on 23/03/2018

Embase search results

#	Terms	Results
1	heroin*.ti,ab.	17939
2	opiate*.ti,ab.	31806
3	opium*.ti,ab.	2914
4	opioid*.ti,ab.	97282
5	morphine*.ti,ab.	64411
6	exp Diamorphine/	22857
7	exp Opiate/	69546
8	exp heroin dependence/	9183
9	exp opiate addiction/	14690
10	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9	210638
11	exp intravenous drug abuse/	9784
12	injecting risk.ti,ab.	175
13	"IDU\$1".tw.	7487
14	"IVDU\$1".tw.	1189
15	"PWID\$1".tw.	1307
16	"injecting drug".tw.	4753
17	"intravenous drug".tw.	9605
18	"injecting substance".tw.	10
19	"intravenous substance".tw.	72
20	exp substance abuse, intravenous/	49527
21	11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19 or 20	72604
22	10 or 21	273105
23	exp intervention study/	35430
24	exp early intervention/	19363

#	Terms	Results
25	drug dependence/di, dm, dt, pc, rh, th [Diagnosis, Disease Management, Drug Therapy, Prevention, Rehabilitation, Therapy]	7797
26	exp drug dependence treatment/	19705
27	exp residential care/	11221
28	exp harm reduction/	4534
29	Addiction/dt [Drug Therapy]	1413
30	23 or 24 or 25 or 26 or 27 or 28 or 29	95877
31	(brief adj3 intervention).ti,ab.	5654
32	(safe adj3 inject*).ti,ab.	1489
33	(needle adj2 syringe adj5 program).ti,ab.	109
34	exp condom/	18629
35	sexually transmitted disease/di, dm, dt, pc, rh, th [Diagnosis, Disease Management, Drug Therapy, Prevention, Rehabilitation, Therapy]	12456
36	hepatitis C/di, dm, dt, pc, rh, th [Diagnosis, Disease Management, Drug Therapy, Prevention, Rehabilitation, Therapy]	38575
37	Human immunodeficiency virus/dm, pc [Disease Management, Prevention]	7
38	exp pre-exposure prophylaxis/	1660
39	(prep adj5 HIV).ti,ab.	996
40	(prep adj5 sexual*).ti,ab.	140
41	(pre-exposure prophylaxis adj5 sexual*).mp. [mp=title, abstract, heading word, drug trade name, original title, device manufacturer, drug manufacturer, device trade name, keyword, floating subheading word]	76
42	(drug adj3 consum* adj3 (room or facil*)).ti,ab.	35
43	(supervis* adj3 inject*).ti,ab.	351
44	atypical antipsychotic agent/ct, dt [Clinical Trial, Drug Therapy]	7690
45	Suicide/pc	5910
46	(suicide adj3 prevent*).ti,ab.	5886
47	exp behavior therapy/	42863

#	Terms	Results
48	exp motivational interviewing/	3338
49	exp cognitive behavioural therapy/	5043
50	exp self help/	12974
51	((peer or mutual) adj3 support*).ti,ab.	6276
52	exp family therapy/	12896
53	multisystemic therapy.ti,ab.	200
54	exp therapeutic community/	3051
55	agonist pharmacotherap*.ti,ab.	32
56	(drug adj3 (compulsory or detention)).ti,ab.	136
57	"NSP\$1".tw.	5311
58	"needle syringe program\$".tw.	75
59	"NSEP\$1".tw.	81
60	"needle syringe exchange program\$".tw.	33
61	"needle exchange\$1".tw.	1078
62	"syringe exchange\$1".tw.	725
63	exp preventive health service/	26240
64	"HAART".tw.	15971
65	"Anti?Retroviral Treatment".tw.	8878
66	"Anti?Retroviral Therapy".tw.	43929
67	"highly active anti?retroviral therapy".tw.	12656
68	"highly active anti?retroviral treatment".tw.	628
69	"HIV treatment".tw.	4851
70	"anti-HIV Agents".tw.	934
71	"AIDS treatment".tw.	1029
72	exp anti human immunodeficiency virus agent/	142530
73	exp human immunodeficiency virus fusion inhibitor/	12218

#	Terms	Results
74	exp highly active antiretroviral therapy/	35463
75	"contingency management".tw.	1075
76	Vaccine/dt [Drug Therapy]	5049
77	detoxification/	23238
78	exp drug detoxification/	4363
79	exp psychosocial care/	15876
80	exp residential care/	11221
81	exp rehabilitation center/	13384
82	exp cognitive therapy/	43099
83	exp "acceptance and commitment therapy"/	851
84	"OST".tw.	1946
85	"opioid substitution".tw.	788
86	"methadone".tw.	16914
87	"MMT".tw.	3538
88	"methadone maintenance".tw.	4772
89	"buprenorphine".tw.	7659
90	"Buprenorphine maintenance".tw.	337
91	"opioid replacement".tw.	155
92	"opioid agonist".tw.	3256
93	"opiate agonist".tw.	604
94	"opioid maintenance".tw.	497
95	"opiate maintenance".tw.	143
96	exp Buprenorphine/	14674
97	exp Methadone/	30108
98	exp Opiate Substitution Treatment/	1612
99	exp methadone treatment/	4162

#	Terms	Results
100	exp buprenorphine plus naloxone/	1297
101	"naltrexone".tw.	7969
102	exp Naltrexone/	13328
103	31 or 32 or 33 or 34 or 35 or 36 or 37 or 38 or 39 or 40 or 41 or 42 or 43 or 44 or 45 or 46 or 47 or 48 or 49 or 50 or 51 or 52 or 53 or 54 or 55 or 56 or 57 or 58 or 59 or 60 or 61 or 62 or 63 or 64 or 65 or 66 or 67 or 68 or 69 or 70 or 71 or 72 or 73 or 74 or 75 or 76 or 77 or 78 or 79 or 80 or 81 or 82 or 83 or 84 or 85 or 86 or 87 or 88 or 89 or 90 or 91 or 92 or 93 or 94 or 95 or 96 or 97 or 98 or 99 or 100 or 101 or 102	531556
104	30 or 103	598139
105	exp evaluation study/	46728
106	exp program evaluation/	15815
107	exp evidence based medicine/	956745
108	exp outcome assessment/	411185
109	exp randomized controlled trial/	491232
110	random* controlled trial.ti,ab.	96486
111	"clinical trial (topic)"/	91565
112	exp controlled clinical trial/	669443
113	multicenter study/	177890
114	random*.ti,ab.	1287665
115	(pretest or pre test).ti,ab.	21706
116	(posttest or post test).ti,ab.	22682
117	before after.ti,ab.	6543
118	qua?irandomi*.ti,ab.	115
119	nonrandomi*.ti,ab.	12767
120	exp open study/	30572
121	exp cohort analysis/	350020
122	exp longitudinal study/	110258

#	Terms	Results
123	systematic review/	161232
124	"meta analysis (topic)"/	36687
125	105 or 106 or 107 or 108 or 109 or 110 or 111 or 112 or 113 or 114 or 115 or 116 or 117 or 118 or 119 or 120 or 121 or 122 or 123 or 124	2801496
126	22 and 104 and 125	11974
127	limit 126 to (human and yr="2000 -Current" and "review")	1823

Searches conducted on 21/03/2018

PsycINFO search results

#	Terms	Results
1	heroin*.ti,ab.	8779
2	opiate*.ti,ab.	8566
3	opium*.ti,ab.	592
4	opioid*.ti,ab.	19819
5	morphine*.ti,ab.	9991
6	exp Heroin addiction/	2489
7	exp Opiates/	22956
8	exp Heroin/	2556
9	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8	39477
10	exp Intravenous Drug Usage/	3669
11	injecting risk.ti,ab.	107
12	"IDU\$1".tw.	2123
13	"IVDU\$1".tw.	119
14	"PWID\$1".tw.	582
15	"injecting drug".tw.	1789
16	"intravenous drug".tw.	1022
17	"intravenous substance".tw.	13

#	Terms	Results
18	"injecting substance".tw.	5
19	10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18	5750
20	9 or 19	43797
21	exp intervention/	91049
22	exp early intervention/	9959
23	exp drug dependency/	24742
24	exp drug therapy/	134549
25	"drug dependence treatment".ti,ab.	87
26	exp harm reduction/	3131
27	exp Addiction/	56195
28	21 or 22 or 23 or 24 or 25 or 26 or 27	285287
29	(brief adj3 intervention).ti,ab.	3983
30	(safe adj3 inject*).ti,ab.	103
31	(needle adj2 syringe adj5 program).ti,ab.	66
32	exp condom/	3696
33	exp sexually transmitted disease/	42686
34	("Hepatitis C" or "HCV").ti,ab.	2737
35	exp HIV/	39644
36	(prep adj5 HIV).ti,ab.	221
37	(prep adj5 sexual*).ti,ab.	41
38	(pre-exposure prophylaxis adj5 sexual*).mp. [mp=title, abstract, heading word, table of contents, key concepts, original title, tests & measures]	8
39	(drug adj3 consum* adj3 (room or facil*)).ti,ab.	22
40	(supervis* adj3 inject*).ti,ab.	167
41	"atypical antipsychotic agent".ti,ab.	166
42	exp Suicide/	25670

#	Terms	Results
43	(suicide adj3 prevent*).ti,ab.	5599
44	exp behavior therapy/	18951
45	exp motivational interviewing/	2131
46	exp cognitive behavior therapy/	18032
47	exp self-help techniques/	9862
48	exp support groups/	5709
49	((peer or mutual) adj3 support*).ti,ab.	5910
50	exp family therapy/	21148
51	multisystemic therapy.ti,ab.	408
52	exp therapeutic community/	2741
53	agonist pharmacotherap*.ti,ab.	20
54	(drug adj3 (compulsory or detention)).ti,ab.	80
55	"NSP\$1".tw.	359
56	"needle syringe program\$".tw.	68
57	"NSEP\$1".tw.	17
58	"needle syringe exchange program\$".tw.	25
59	"needle exchange\$1".tw.	509
60	"syringe exchange\$1".tw.	404
61	exp preventive medicine/	2017
62	"HAART".tw.	1250
63	"Anti?Retroviral Treatment".tw.	954
64	"Anti?Retroviral Treatment".tw.	954
65	"Anti?Retroviral Therapy".tw.	4142
66	"highly active anti?retroviral therapy".tw.	1085
67	"highly active anti?retroviral treatment".tw.	68
68	"HIV treatment".tw.	1067

#	Terms	Results
69	"anti-HIV Agents".tw.	5
70	"AIDS treatment".tw.	248
71	"anti human immunodeficiency virus agent".ti,ab.	0
72	"human immunodeficiency virus fusion inhibitor".ti,ab.	0
73	"contingency management".tw.	1605
74	exp Immunization/	4076
75	detoxification/	1683
76	exp drug rehabilitation/	28677
77	exp psychosocial rehabilitation/	10810
78	exp rehabilitation centers/	1026
79	exp cognitive therapy/	12909
80	exp "acceptance and commitment therapy"/	1429
81	"OST".tw.	467
82	"opioid substitution".tw.	408
83	"methadone".tw.	7082
84	"MMT".tw.	786
85	"methadone maintenance".tw.	3421
86	"buprenorphine".tw.	2399
87	"Buprenorphine maintenance".tw.	213
88	"opioid replacement".tw.	58
89	"opioid agonist".tw.	951
90	"opiate agonist".tw.	142
91	"opioid maintenance".tw.	254
92	"opiate maintenance".tw.	57
93	exp Buprenorphine/	1588
94	exp Methadone/	1740

#	Terms	Results
95	"opiate substitution".ti,ab.	115
96	exp methadone maintenance/	3428
97	(buprenorphine and naloxone).ti,ab.	448
98	exp naltrexone/	1789
99	"naltrexone".tw.	3208
100	29 or 30 or 31 or 32 or 33 or 34 or 35 or 36 or 37 or 38 or 39 or 40 or 41 or 42 or 43 or 44 or 45 or 46 or 47 or 48 or 49 or 50 or 51 or 52 or 53 or 54 or 55 or 56 or 57 or 58 or 59 or 60 or 61 or 62 or 63 or 64 or 65 or 66 or 67 or 68 or 69 or 70 or 71 or 72 or 73 or 74 or 75 or 76 or 77 or 78 or 79 or 80 or 81 or 82 or 83 or 84 or 85 or 86 or 87 or 88 or 89 or 90 or 91 or 92 or 93 or 94 or 95 or 96 or 97 or 98 or 99	217256
101	28 or 100	454618
102	exp treatment effectiveness evaluation/	22349
103	exp program evaluation/	19270
104	exp evidence based practice/	15874
105	"outcome assessment".ti,ab.	870
106	random* controlled trial.ti,ab.	16786
107	exp clinical trials/	10840
108	"controlled clinical trial".ti,ab.	1312
109	"multicenter study".ti,ab.	1362
110	random*.ti,ab.	175803
111	(pretest or pre test).ti,ab.	16361
112	(posttest or post test).ti,ab.	22005
113	before after.ti,ab.	820
114	qua?irandomi*.ti,ab.	12
115	nonrandomi*.ti,ab.	881
116	exp cohort analysis/	1248
117	exp longitudinal studies/	15918

#	Terms	Results
118	"systematic review".ti,ab.	19740
119	exp meta analysis/	4150
120	102 or 103 or 104 or 105 or 106 or 107 or 108 or 109 or 110 or 111 or 112 or 113 or 114 or 115 or 116 or 117 or 118 or 119	281942
121	20 and 101 and 120	2986
122	limit 121 to (human and "reviews (best balance of sensitivity and specificity)" and yr="2000 -Current")	1669

Searches conducted on 27/03/2018

Table 3. Grades of evidence

Grade	Evidence
A	Consistent conclusions across meta-analyses, high quality systematic reviews of randomised controlled trials, or several randomised controlled trials
B	Evidence from one or two randomised controlled trials only
C	High-quality systematic reviews with some inconsistent conclusions from authors, or several consistent ecological studies or cohort studies
D	Cross-sectional association, case series suggesting outcome, or a single cohort study

Description of evidence on reviews of interventions to reduce health outcomes

Opioid Agonist Treatment (OAT)

Outcome examined	First author	Year	Impact	Effect	95% CI	95% CI	Measure	Level of evidence	Number of studies in review	Notes	Author notes on quality of evidence
All-cause mortality	Mattick	2014	?	0.48	0.1	2.39	RR	Systematic review and MA of RCTs	4		Moderate quality evidence
All-cause mortality	Sordo	2017	↓					Systematic review and MA of cohort studies	19		noted potential for confounding and selection bias as all cohorts
All-cause mortality	Crowley	2017	↓					Systematic review of RCTs and other study designs	Eight randomised control trials (RCT), one non-randomised trial and five observational studies	taken from abstract, could not access paper	
All-cause mortality	Ayanga	2016	?					Review (methods unclear)		referred to Mattick et al Cochrane reviews	
Contact with the criminal justice system	Perry	2015	×	0.6	0.32	1.14	RR	Systematic review and MA of RCTs	1	arrest	Low quality evidence
Contact with the criminal justice system	Perry	2015	×	0.77	0.36	1.64	RR	Systematic review and MA of RCTs	3	reincarceration	Low quality evidence
Contact with the criminal justice system	Maglione	2018	×	0.75	0.46	1.23	RR	Systematic review and MA of RCTs and cohort studies	6	arrest or incarceration	
Criminal activity	Mattick	2014	×	0.39	0.12	1.25	RR	Systematic review and MA of RCTs	3	any criminal activity	Moderate quality evidence
Criminal activity	Ayanga	2016	?					Review (methods unclear)		referred to Mattick et al Cochrane reviews	
Criminal activity	Perry	2015	↓	-74.2	133.5	14.89	MD	Systematic review and MA of RCTs	1	criminal activity continuous	Low quality evidence
Criminal activity	Maglione	2018	↓	-0.57	-1.00	-0.13	SMD	Systematic review and MA of RCTs and cohort studies	2	criminal activity continuous	
HCV incidence	Connery	2015	↓					Systematic review of RCTs and cohort studies	unclear		
HCV incidence	Hagan	2012	×	0.60	0.35	1.03	RR	Systematic review and MA of cohort studies	8		No information on quality of evidence
HCV incidence	MacArthur	2014	↓					Review of reviews			There is tentative review-level evidence to support the effectiveness of OST in reducing HCV transmission
HCV incidence	Platt	2017	↓	0.50	0.40	0.63	RR	Systematic review and MA of cohort studies	21		Low quality evidence
HIV incidence	Gowing	2011	↓					Systematic review and MA of RCTs and cohort studies	4		
HIV incidence	Connery	2015	↓					Systematic review of RCTs and cohort studies	unclear		
HIV incidence	MacArthur	2012	↓	0.46	0.32	0.67	RR	Systematic review and MA of cohort studies	9		
HIV incidence	MacArthur	2014	↓					Review of reviews			There is sufficient review-level evidence to conclude that OST is effective in reducing HIV seroconversion
HIV linkage to care and treatment	Low	2016	↑	1.54	1.17	2.03	OR	Systematic review and MA of cohort studies	10	Odds of being on HIV therapy	
HIV linkage to care and treatment	Low	2016	↑	1.87	1.50	2.33	HR	Systematic review and MA of cohort studies	4	Effect of OST on initiating HIV therapy	
HIV treatment adherence	Low	2016	↑	2.14	1.41	3.26	OR	Systematic review and MA of cohort studies	5	Effect of OST on adherence to HIV therapy	
HIV viral suppression	Low	2016	↑	1.45	1.21	1.73	OR	Systematic review and MA of cohort studies	10	Effect of OST on HIV viral load suppression	
Injecting frequency	Gowing	2011	↓	-0.59	-0.91	-0.26	SMD	Systematic review and MA of RCTs and cohort studies	10	we have pooled estimates - Gowing did not - RCTs	
Injecting risk behaviour	Gowing	2011	↓	0.53	0.4	0.7	RR	Systematic review and MA of RCTs and cohort studies	12	we have pooled estimates - Gowing did not - RCTs	
Injecting risk behaviour	Connery	2015	↓					Systematic review of RCTs and cohort studies	unclear		
Injecting risk behaviour	Macarthur	2014	↓					Review of reviews			In light of primary evidence and statements of evidence from core reviews in support of OST, we concluded that there is sufficient evidence to support effectiveness of OST in reducing IRB.
Mental health	Fingleton	2015	↑					Systematic review of RCTs and cohort studies	19 RCTs and 3 cohorts	varied assessment of mental health across studies	
Mental health	Maglione	2018	×					systematic review	3	differing measurements; ASI and SF-12	
Mental health	Feelmeyer	2014	↑	0.491	0.351	0.631	SMD	Systematic review and MA of cohort studies	8 with 9 samples	Clients in OST in low and middle income countries; used WHOQOL-BREF and ASI - reported separately by different domains; this one is psychological	
Neonatal outcomes	Tran	2017						Systematic review of RCTs and other study designs	3 RCTs and 8 observational studies	birth weight - all methadone vs. bupe	
Neonatal outcomes	Tran	2017						Systematic review of RCTs and other study designs	3 RCTs and 8 observational studies	neonatal abstinence syndrome - all methadone vs. bupe	
Neonatal outcomes	Minozzi	2013	↑	-365	-673	-57	MD	Systematic review and MA of RCTs	2	birth weight	Low quality evidence
Neonatal outcomes	Minozzi	2013	×	1.22	0.89	1.67	RR	Systematic review and MA of RCTs	3	number treated for neonatal abstinence	Very low quality evidence
Neonatal outcomes	Minozzi	2013	×	0	-0.03	0.03	MD	Systematic review and MA of RCTs	2	APGAR score	Very low quality evidence

Outcome examined	First author	Year	Impact	Effect	95% CI	95% CI	Measure	Level of evidence	Number of studies in review	Notes	Author notes on quality of evidence
Neonatal outcomes	Hand	2017						Review (methods unclear)			
Neonatal outcomes	Zedler	2016	↑	0.40	0.18	0.91	RR	Systematic review and MA of RCTs and cohort studies	This outcome had 2 RCTs; overall review had Three RCTs (n = 223) and 15 cohort OBSs (n = 1923) - have extracted RCT only	pre-term birth	Low quality evidence
Neonatal outcomes	Zedler	2016	↑	324	32	617	RCT WMD	Systematic review and MA of RCTs and cohort studies	This outcome had 2 RCTs; overall review had Three RCTs (n = 223) and 15 cohort OBSs (n = 1923) - have extracted RCT only	birth weight	Low quality evidence
Neonatal outcomes	Zedler	2016	↑	0.90	0.14	1.66	RCT WMD	Systematic review and MA of RCTs and cohort studies	This outcome had 2 RCTs; overall review had Three RCTs (n = 223) and 15 cohort OBSs (n = 1923) - have extracted RCT only	head circumference	Low quality evidence
Neonatal outcomes	Noormohammadi	2016						Systematic review of RCTs and other study designs	5 studies		
Neonatal outcomes	Holbrook	2015						Review of reviews	9 reviews and one MA		
Opioid use	Mattick	2014	↓	0.66	0.56	0.78	RR	Systematic review and MA of RCTs	6	morphine positive urine or hair	High quality evidence
Opioid use	Mattick	2014	↓				RR	Systematic review and MA of RCTs	6	self-reported heroin use	High quality evidence
Opioid use	Nielsen	2016	↓	0.63	0.43	0.91	RR	Systematic review and MA of RCTs	3	morphine positive urine or hair	Low quality evidence
Opioid use	Nielsen	2016	↓	0.54	0.31	0.93	RR	Systematic review and MA of RCTs	3	self-reported opioid use	Low quality evidence
Opioid use	Gowing	2011	↓	0.48	0.41	0.55	RR	Systematic review and MA of RCTs and cohort studies	11	we have pooled estimates from follow-up studies	
Opioid use	Ayanga	2016	↓					Review (methods unclear)		referred to Mattick et al Cochrane reviews	
Opioid use	Thomas	2015	↓					Systematic review of reviews, MAs, RCTs and cohort studies	7 reviews or MAs, 16 RCTs, a randomized cross-over study, a study using a self-administered survey, and a retrospective descriptive study		High quality evidence
Overdose mortality	Sordo	2017	↓					Systematic review and MA of cohort studies	19		noted potential for confounding and selection bias as all cohorts
Overdose mortality	Connery	2015	↓					Systematic review of RCTs and cohort studies	unclear		
Overdose mortality	Ma	2018	↓	0.12	0.06	0.22	RR	Systematic review and MA of cohort studies	6	in treatment vs. untreated	
Overdose mortality	Ma	2018	↓	0.32	0.25	0.42	RR	Systematic review and MA of cohort studies	14	in treatment versus out of treatment	
Quality of life	Fei	2016	↑					Retrospective cohort		Cohort of patients retained in treatment in Malaysia	
Quality of life	Feelmeyer	2014	↑	0.292	0.164	0.421	SMD	Systematic review and MA of cohort studies	8 with 9 samples	Clients in OST in low and middle income countries; used WHOQOL-BREF and ASI - reported separately by different domains; this one is social	
Quality of life	Maglione	2018	↑								
Quality of life	Thornton	2017	X	-0.05	-0.18	0.08	Hedge's G effect size	Systematic review	3	people receiving long term opioid therapy for pain - mental component summary score	
Quality of life	Thornton	2017	↑	0.18	0.08	0.28	Hedge's G effect size	Systematic review	3	people receiving long term opioid therapy for pain - physical component summary score	
Skin and soft tissue infections	Larney	2017	?					Systematic review	3	all cross-sectional studies assessing whether past year or ever OST associated with past 6-12 month abscesses	

OAT vs. no treatment

Overdose

Table J1. Calculated overdose deaths rate ratios using data identified in Sordo, 2017 and Hickman, 2019

Study	Rate ratio	95% confidence interval		% Weight
		Lower	Upper	
Buster, 2002	0.946	0.567	1.608	8.92
Caplehorn, 1996	0.235	0.058	0.708	4.75
Clausen, 2008	0.173	0.097	0.310	8.51
Cousins, 2016	0.601	0.377	1.050	8.97
Cushman, 1977	0.103	0.022	0.403	3.96
Davoli, 2007	0.135	0.043	0.407	5.31
Gearing, 1974	0.127	0.071	0.231	8.49
Grönbladh, 1990	0.177	0.065	0.416	6.36
Hickman, 2018 (methadone)	0.422	0.252	0.699	8.99
Hickman, 2018 (buprenorphine)	0.319	0.090	0.938	5.10
Kimber, 2015 (methadone)	0.386	0.314	0.474	10.56
Kimber, 2015 (buprenorphine)	0.309	0.202	0.458	9.60
Peles, 2010	0.070	0.020	0.210	5.04
Scherbaum, 2002	0.071	0.022	0.201	5.44
Pooled rate ratio	0.253	0.176	0.363	100.00

Overdose Mortality Rate Ratio




Figure J1. Pooled rate ratio of overdose deaths within buprenorphine/methadone treatment vs. no treatment (ref) as calculated using data within Sordo, 2017 and Hickman, 2018.

All-cause mortality

Table J2. Calculated all-cause mortality rate ratios using data identified in Sordo, 2017

Study	Rate ratio	95% confidence interval		% Weight
		Lower	Upper	
Caplehorn, 1996	0.353	0.162	0.708	3.05
Clausen, 2008	0.395	0.274	0.577	5.67
Cornish, 2010 (methadone)	0.353	0.222	0.548	4.98
Cornish, 2010 (buprenorphine)	0.710	0.230	2.068	1.74
Cousins, 2016	0.324	0.245	0.428	6.51
Cushman, 1977	0.320	0.160	0.667	3.18
Degenhardt, 2009	0.407	0.370	0.446	7.82
Evans, 2015	0.366	0.308	0.433	7.38
Fugelstad, 1995	0.298	0.086	1.156	1.32
Fugelstad, 1998	0.564	0.143	2.625	1.09
Fugelstad, 2007	0.407	0.292	0.567	6.03
Gearing, 1974	0.269	0.181	0.411	5.34
Grönbladh, 1990	0.341	0.175	0.640	3.55
Hickman, 2018 (methadone)	0.357	0.212	0.585	7.30
Hickman, 2018 (buprenorphine)	0.305	0.254	0.366	4.53
Kimber, 2015 (methadone)	0.557	0.497	0.625	7.72
Kimber, 2015 (buprenorphine)	0.395	0.308	0.502	6.80
Nosyk, 2015	0.172	0.132	0.221	6.70
Peles, 2010	0.147	0.096	0.226	5.18
Reece, 2010	0.463	0.092	1.454	1.19
Scherbaum, 2002	0.199	0.093	0.431	2.91
Pooled rate ratio	0.334	0.284	0.394	100.00

All Cause Mortality Rate Ratio




Figure J2. Pooled rate ratio of all-cause mortality within buprenorphine/methadone treatment vs. no treatment (ref) using calculated rate ratios from data within Sordo, 2017 and Hickman, 2018.

Systematic reviews of OAT versus no treatment in prison

Outcome examined	First author	Year	Impact	Effect	95 % CI	95 % CI	Author notes on quality of evidence
Opioid use	Hedrich	2012	↓				
Injecting risk behaviour	Hedrich	2012					
HIV incidence	Hedrich	2012	?				insufficient evidence
HCV incidence	Hedrich	2012	?				insufficient evidence
Opioid use	Larney	2010	↓				
Injecting risk behaviour	Larney	2010	↓				
HIV incidence	Larney	2010	?				no evidence

Buprenorphine vs. methadone (ref)

Overdose during induction (4 weeks)

Table J3. Overdose deaths rate ratios within first 4 weeks of treatment

Study	Rate ratio	95% confidence interval		% Weight
		Lower	Upper	
Hickman, 2018	0.240	0.030	1.960	42.78
Kimber, 2015	0.177	0.020	0.741	57.22
Pooled rate ratio	0.202	0.051	0.792	100.00




Figure J3. Pooled rate ratio of overdose deaths within first 4 weeks of treatment.

Overdose following cessation (4 weeks)

Table J4. Overdose death rate ratios within first 4 weeks out of treatment

Study	Rate ratio	95% confidence interval		% Weight
		Lower	Upper	
Hickman, 2018	1.170	0.460	2.970	45.73
Kimber, 2015	1.974	0.864	4.787	54.27
Pooled rate ratio	1.554	0.827	2.920	100.00




Figure J4. Pooled rate ratio of overdose deaths within first 4 weeks out of treatment.

Panel J1 – Other opioid agonist treatments (OAT) for opioid dependence

Injectable opioids are prescribed in a small number of countries for the treatment of opioid dependence refractory to oral opioid substitution treatment. Diamorphine (pharmaceutical heroin) is the most widely prescribed injectable opioid, currently used to treat opioid dependence in the United Kingdom, Denmark, Germany, the Netherlands, Switzerland, and Canada. Injectable doses of diamorphine are taken under direct medical or nursing supervision.

A recent systematic review of supervised injectable diamorphine for the treatment of opioid dependence identified six randomised controlled trials.⁶² The comparator in each trial was oral methadone, with some studies including an additional comparator. A meta-analysis suggested significantly better retention in injectable diamorphine compared to oral methadone (RR 1.37; 95%CI 1.03–1.83). Extra-medical opioid use was measured differently across trials, precluding meta-analysis, but all five of the trials to examine this outcome reported a significantly greater suppression of illicit heroin use with injectable diamorphine compared to oral methadone.

The Randomised Injectable Opiate Treatment Trial (RIOTT) was a three-arm randomised controlled trial comparing injectable diamorphine, injectable methadone, and optimised oral methadone (daily doses of >80mg with psychosocial support). Injectable diamorphine was superior to injectable and oral methadone at suppressing illicit heroin use; injectable methadone was not superior to oral methadone in suppressing illicit heroin use.⁶³

Injectable hydromorphone was compared to injectable diamorphine in the Study to Assess Longer-term Opioid Medication Effectiveness (SALOME) study. Non-inferiority of injectable hydromorphone to injectable diamorphine in suppressing illicit opioid use was confirmed.⁶⁴ Injectable hydromorphone is offered alongside injectable diamorphine as a treatment option in Vancouver, British Columbia.⁶⁵

References

1. Strang J, Groshkova T, Uchtenhagen A, et al. Heroin on trial: systematic review and meta-analysis of randomised trials of diamorphine-prescribing as treatment for refractory heroin addiction. *Br J Psychiatry* 2015; **207**(1): 5-14.
2. Strang J, Metrebian N, Lintzeris N, et al. Supervised injectable heroin or injectable methadone versus optimised oral methadone as treatment for chronic heroin addicts in England after persistent failure in orthodox treatment (RIOTT): a randomised trial. *Lancet (London, England)* 2010; **375**(9729): 1885-95.
3. Oviedo-Joekes E, Guh D, Brissette S, et al. Hydromorphone compared with diacetylmorphine for long-term opioid dependence. *JAMA psychiatry* 2016; **73**(5): 447-55.
4. Tyndall M. An emergency response to the opioid overdose crisis in Canada: a regulated opioid distribution program. *Canadian Medical Association Journal* 2018; **15**: E35-6.

Panel J2 – Improving opioid agonist treatment access, retention and outcomes

There are many opportunities to improve the quality of care provided in OAT. Key barriers to treatment entry include: lack of unsupervised dosing, extensive assessment processes, delays to treatment following presentation, mandatory engagement in psychosocial counselling, siloing of OAT in specialist clinics, and geographically distant clinics and pharmacies for medication delivery.

In many countries, there is a requirement for official registration of a patient due to the scheduling of opioid medications. This information can sometimes be shared with police (either by law, or in practice), and clients may be targeted by police, lose their driver's license, experience restrictions on employment, and other adverse consequences^{1,2}.

Opportunities to improve OAT access, retention and outcomes:

- Rapid access to OAT with minimal assessment to establish opioid dependence in several settings such as emergency rooms, community-based settings, primary care and criminal justice settings
- OAT prescribing in a range of settings, beyond specialised settings including primary care, and OAT dispensing in community pharmacies
- Elimination of compulsory counselling requirements (e.g. in the US); instead, make provision of psychosocial services that may be accessed on a voluntary basis
- Minimise intensity of care as people stabilise in treatment (e.g., transition to less supervision as patients become clinically stable)
- Increased and rapid access to unsupervised dosing, particularly for people prescribed buprenorphine or stabilised on methadone
- Minimise use of urine drug screening; use results of such screening to improve care (i.e. dose adjustment) rather than to terminate treatment
- Low-threshold and adapted care for disengaged individuals, such as those who are homeless or with mental health issues (e.g., many required assessments can be obtained after starting OAT)
- Access to alternative agonist treatments such as slow release oral morphine and injectable opioid agonist therapy for individuals who do not respond to methadone or buprenorphine

References

1. Bojko MJ, Mazhnaya A, Makarenko I, et al. "Bureaucracy & Beliefs": Assessing the barriers to accessing opioid substitution therapy by people who inject drugs in Ukraine. *Drug-Educ Prev Polic* 2015; **22**(3): 255-62.
2. Bojko MJ, Dvoriak S, Altice FL. At the crossroads: HIV prevention and treatment for people who inject drugs in Ukraine. *Addiction* 2013; **108**(10): 1697-9.

Provision of sterile injecting equipment

Outcome examined	First author	Year	Impact	Effect	95% CI	95% CI	Measure	Level of evidence	Number of studies in review	Notes	Author notes on quality
HCV incidence	Fernandes	2017						Review of reviews			
HCV incidence	Hagan	2012	↑	1.62	1.04	2.52	RR	Systematic review and MA of cohort studies	7	Volunteer bias in studies which found an association between HCV acquisition and PWID	
HCV incidence	Turner	2011	↓	0.48	0.25	0.93	aOR	Individual participant data meta-analysis	3	Evaluated high NSP coverage (>=100%) compared to <100%	
HCV incidence	MacArthur	2014	?					Review of reviews			Insufficient evidence
HCV incidence	Platt	2017	x	0.77	0.38	1.54	RR	Systematic review and MA of cohort studies	10	Overall (NA + Europe)	
HCV incidence	Platt	2017	↓	0.44	0.24	0.80	RR	Systematic review and MA of cohort studies	6	Europe alone	
HCV incidence	Platt	2017	x	1.58	0.57	4.42	RR	Systematic review and MA of cohort studies	3	NA alone	
HCV incidence	Sawangjit	2017	?					Systematic review and meta-analysis		review of pharmacy-based NSP	Insufficient evidence
HIV incidence	Fernandes	2017	↓					Review of reviews			
HIV incidence	Aspinall	2014	↓	0.66	0.43	1.01	OR/HR/RR	Systematic review and MA of cohort studies	12	Across all studies	
HIV incidence	Aspinall	2014	↓	0.42	0.22	0.81	OR/HR/RR	Systematic review and MA of cohort studies	6	Higher quality studies (according to Newcastle-Ottawa tool)	
HIV incidence	MacArthur	2014	↓					Review of reviews			
HIV incidence	Sawangjit	2017	?					Systematic review and meta-analysis		review of pharmacy-based NSP	Insufficient evidence
Injecting risk behaviour	Turner	2011	↓	0.52	0.32	0.83	aOR	Individual participant data meta-analysis		Evaluated high NSP coverage (>=100%)	
Injecting risk behaviour	Fernandes	2017						Review of reviews			
Injecting risk behaviour	MacArthur	2014	↓					Review of reviews	43		Sufficient evidence
Injecting risk behaviour	Sawangjit	2017	↓	0.5	0.34	0.73	OR	Systematic review and meta-analysis		review of pharmacy-based NSP	
Skin and soft tissue infections	Larney	2017	?					Systematic review	3	all cross-sectional studies assessing whether past year or current NSP associated with reduced risk of skin and soft tissue infections	

Naloxone

Outcome examined	First author	Year	Impact	Effect	95% CI	95% CI	Measure	Level of evidence	Number of studies in review	Notes
Non-fatal overdose	McAuley	2015	?	9.2	5.2	13.1	users per 3 months per 100 PWID		25	uses per 100 PWID trained and given naloxone; estimated 40 uses per 100 PWID trained per year, but 9.2*4 = 37
Non-fatal overdose	Clark	2014	?	-						
	Mueller	2015	↓	-						examined cohort studies

Drug consumption rooms (DCRs)

Outcome examined	First author	Year	Impact	Effect	95% CI	95% CI	Measure	Level of evidence	Number of studies in review	Notes	Author notes on quality of evidence
Non-fatal overdose	Potier	2014	↓					Systematic review that located a simulation study, ITS, cross-sectional survey	3		
Overdose mortality	Potier	2014	↓					Systematic review that located an ITS study	1		
Injecting risk behaviour	Potier	2014	↓					Systematic review that located a MA, a cross-sectional survey	4		
Skin and soft tissue infections	Potier	2014	?				incidence	Systematic review that located three papers on the SEOSI cohort	3	3 papers examining SEOSI cohort, mentions no evidence for 'viral transmission ' in the same paragraph. Is it worth doing a '?' for each of HIV incidence, HICv incidence, etc.	
Injecting risk behaviour	Milloy	2009	↓	0.31	0.17	0.55	RR	MA of three cohorts	3	Look at sharing, lending and borrowing needles. Sep. info for HIV negative and positive	
Skin and soft tissue infections	Larney	2017	↓	0.47	0.23	0.94	OR	Systematic review	1	1 cross-sectional study, Unadjusted, 'Current-1 month'	
Injecting risk behaviour	Macarthur	2014	↓					Review of reviews		statement of insufficient evidence from a core review, tentative statements of evidence from two supplementary reviews, and evidence from a number of robust studies	Tentative review-level evidence that injecting risks decrease
HIV incidence	Macarthur	2014	?					Review of reviews	1 cross-sectional study identified across reviews		Insufficient evidence to either support or discount the effectiveness of SIFs in preventing HIV
HCV incidence	Macarthur	2014	?					Review of reviews	1 cross-sectional study identified across reviews		Insufficient evidence to either support or discount the effectiveness of SIFs in preventing HCV

Oral opioid antagonists

Outcome examined	First author	Year	Journal	Title	Impact	Effect	95 % CI	95 % CI	Measure	Level of evidence	Number of studies in review	Notes
Contact with the criminal justice system	Maglione	2018	Journal of Substance Abuse Treatment	Effects of medication assisted treatment (MAT) for opioid use disorder on functional outcomes: A systematic review	x				MD	systematic review	1	number of charges or number of arrests reported
Opioid use	Jahagirdar, D.	2017	CADTH evidence review	Naltrexone for Opioid Use Disorders: A Review of Clinical Effectiveness, Cost-Effectiveness, and Guidelines	X					systematic review		Ten randomized controlled trials ¹³⁻²² and four systematic reviews were included in the review. ²³⁻²⁶ Two of the trials were pilot/proof-of-concept studies. ^{18,19} One study ¹⁴ was a SUMMARY WITH CRITICAL APPRAISAL Naltrexone for Opioid Use Disorders 6 secondary analysis of data from a trial that was also included in this report. ¹³ Three of the systematic reviews included both randomized controlled trials and observational

Opioid
use

Minozzi	2011	John Wiley & Sons, Ltd	Oral naltrexone maintenance treatment for opioid dependence	x
---------	------	------------------------	---	---

1.39 0.6 3.1 abstinenc systemati
1 7 e c review

studies,
and included two,²³
six²⁴ and four²⁵
studies of naltrexone
respectively. The last
systematic
review included
thirteen randomized or
controlled clinical trials
on oral naltrexone.²⁶
The
most recent systematic
review included
publications until
December 2016,²⁴
while the
others included
literature up to
December 2009,²⁵
June 2010,²⁶ and
September 2009.²³

Sustained release opioid antagonists

Outcome examined	First author	Year	Journal	Title	Impact	Effect	95% CI	95% CI	Measure	Level of evidence	Number of studies in review	Notes
Contact with the criminal justice system	Maglione	2018	Journal of Substance Abuse Treatment	Effects of medication assisted treatment (MAT) for opioid use disorder on functional outcomes: A systematic review	↓	-0.23	0.46	-	SMD	Systematic review	1 RCT	mean number of arrest at 78 weeks
Opioid use	Larney	2014	Drug and Alcohol Review	Systematic review of naltrexone implants for opioid dependence	↓	0.57	0.48	0.68	RR	systematic review and MA where possible	5 RCTs	compared to placebo; no difference cf. TAU
Non-fatal overdose	Larney	2014	Drug and Alcohol Review	Systematic review of naltrexone implants for opioid dependence	?					systematic review	2 RCTs, 1 non-RCT	no difference from oral naltrexone or treatment as usual
All-cause mortality	Larney	2014	Drug and Alcohol Review	Systematic review of naltrexone implants for opioid dependence	?					systematic review and MA where possible	4 RCTs, 3 non-RCTs	
Opioid use	Jahagirdar, D.	2017	CADTH evidence review	Naltrexone for Opioid Use Disorders: A Review of Clinical Effectiveness, Cost-Effectiveness, and Guidelines	↓						Ten randomized controlled trials and four systematic reviews	
Non-fatal overdose	Jarvis	2018			↓?					systematic review and MA where possible		no overdoses noted; comparative analysis not possible

Residential rehabilitations

Outcome examined	First author	Year	Impact	Effect	95% CI	95% CI	Measure	Level of evidence	Number of studies in review
Opioid use	Malivert	2012	↓					prospective and retrospective studies	12

HIV testing and informing of serostatus

Outcome examined	First author	Year	Impact
Injecting risk behaviour	Schlumberger	1999	↓

HCV testing and informing of serostatus

Outcome examined	First author	Year	Impact	Effect	95% CI	95% CI	Measure	Level of evidence	Number of studies in review	Notes
Injecting drug use	Spelman	2015	↓	0.95	0.93	0.96	aOR	Merged cohort study analysis	NA	HCV-positive diagnosed PWID demonstrated a 5% per 3-month reduction post-notification in the odds of recent injection drug use
Injecting risk behaviour	Spelman	2015	X	0.97	0.94	1.00	aOR	Merged cohort study analysis	NA	HCV-positive diagnosed PWID demonstrated a 3% per 3-month reduction post-notification in the odds of receptive syringe sharing

HIV treatment

Outcome examined	First author	Year	Journal	Title	Impact	Effect	95% CI	95% CI	Measure	Level of evidence	Number of studies in review	Notes
Injecting risk behaviour	Wood	2012	Current opinion in HIV and AIDS	HIV treatment as prevention among injection drug users	X						3	all cohort studies; no quantitative synthesis and two very old studies
Injecting risk behaviour	Kuyper	2011	Addictive Behaviors	Does initiation of HIV antiretroviral therapy influence patterns of syringe lending among injection drug users?	X	0.87	0.42	1.45	aOR		1	1 cohort study

HCV treatment

Suicide prevention strategies

Outcome examined	First author	Year	Title	Impact	Effect	95% CI	Measure	Level of evidence	Number of studies in review	Notes	Author notes on quality of evidence
Suicide	Haughton	2015	?				7	studies of people with history of self-harm		low to very low evidence; not possible to make a firm conclusion	

Limits on opioid prescribing

n/a

Prescription opioid monitoring programmes

Outcome examined	First author	Year	Journal	Title	Impact	Effect	95% CI	95% CI	Measure	Level of evidence	Number of studies in review	Notes	Author notes on quality of evidence
Non-fatal overdose	Fink	2018	Annals of Internal Medicine	Association between prescription drug monitoring programmes and nonfatal and fatal drug overdoses	?					Systematic review	3	Authors noted serious risk of bias in studies	Low strength evidence - evidence is "insufficient" to make a conclusion
Overdose mortality	Fink	2018	Annals of Internal Medicine	Association between prescription drug monitoring programmes and nonfatal and fatal drug overdoses	?					Systematic review	10	Authors noted serious risk of bias in studies	Low strength evidence - evidence is "insufficient" to make a conclusion

Abuse-deterrent opioids

Outcome examined	First author	Year	Impact	Level of evidence	Number of studies in review	Notes	Author notes on quality of evidence
All-cause mortality	ICER	2017	?	Systematic review	6	all ecological studies; specific drug declined but on consistent trends in overall - unclear if overall impact, other opioids increased	"extremely mixed" "limited evidence" "no consistent trends across studies"
Opioid use	ICER	2017	?	Systematic review	16	pre/post studies of poison centre data (3), treatment entrants (6) and cohorts (3)	evidence is "mixed" for opioid "abuse"

Compulsory drug detention centres

Werb D, Kamarulzaman A, Meacham MC, et al. The effectiveness of compulsory drug treatment: A systematic review. The International journal on drug policy 2016; 28: 1-9.

Criminalisation of drug use

DeBeck K, Cheng T, Montaner JS, et al. HIV and the criminalisation of drug use among people who inject drugs: a systematic review. The Lancet HIV 2017; 4(8): e357-e74.

Table J5: Current evidence for effects of interventions to address key outcomes and behaviours among people who use opioids extra-medically

Intervention	Injecting risk behaviours				Extra-medical opioid use				HIV incidence				HCV incidence				Skin and soft tissue infections				Quality of life			
	Effect	Size of effect	Level	Sources	Effect	Size of effect	Level	Sources	Effect	Size of effect	Level	Sources	Effect	Size of effect	Level	Sources	Effect	Size of effect	Level	Sources	Effect	Size of effect	Level	Sources
Provision of sterile injecting equipment	↓	aOR 0.52 (0.32, 0.83)	A ^{PWID}	1	-	-	-	-	↓	OR/HR/RR 0.42 (0.22, 0.81)	C ^{PWID}	2	↓?	RR 0.77 (0.38, 1.54)	C	3	?	-	D	4	-	-	-	-
Condom provision	-	-	-	-	-	-	-	-	↓	RR 0.29, (0.20, 0.43)	A ^{GEN}	5	?	-	C	6	-	-	-	-	-	-	-	-
Naloxone	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Drug consumption rooms	↓	RR 0.31 (0.17, 0.55)	C ^{PWID}	7	-	-	-	-	?	-	D	8	?	-	D	8	↓	OR 0.47 (0.23, 0.94)	E	4	-	-	-	-
Peer-based self-help groups	-	-	-	-	↓?	-	B ^{ALC}	9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Psychosocial interventions	↓	SMD -0.43 (-0.69, -0.18)	A	10	↓	WMES -0.18 (-0.30, -0.06)	A	11	?	-	D	8	?	-	D	8	?	-	E	12	-	-	-	-
Opioid detoxification alone					✗																			
Oral opioid antagonists	✗	NE	A	13	✗	RR 1.39 (0.61, 3.17)	A	14	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Extended-release opioid antagonists	↓	NE	A	15	↓	NE	A	15,16	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Opioid agonist treatment	↓	RR 0.53 (0.4, 0.7) <small>SYNTH</small>	A	17	↓	RR 0.48 (0.41, 0.55) <small>SYNTH</small>	A	17	↓	RR 0.46 (0.32, 0.67)	C	18	↓	RR 0.50 (0.40, 0.63)	C	3	?	-	D	4	↑	SMD 0.29 (0.16, 0.42)	C	19
Residential rehabilitation	↓	NE	C	20	↓	NE	C	20	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
HIV testing + informing of serostatus	↓	NE	D ^{PWID}	21	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
HCV testing + informing of serostatus	✗	aOR 0.97 (0.94, 1.00)	C ^{PWID}	22	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
HIV treatment	✗	aOR 0.78 (0.42-1.45)	D	23	-	-	-	-	↓	-	D	24	-	-	-	-	-	-	-	-	-	-	-	-
HCV treatment	↓	NE	D ^{PWID}	25	↓	NE	D ^{PWID}	25	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
STI treatment	-	-	-	-	-	-	-	-	↓		A ^{GEN}	26-28	-	-	-	-	-	-	-	-	-	-	-	-
Suicide prevention strategies	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Opioid prescribing limits	-	-	-	-	↓?	NE	D ^{GEN}	29	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Abuse-deterrent opioid formulations	↓	NE	D	30,31	?	NE	D	31	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Prescription opioid monitoring programs	-	-	-	-	↓?	NE	D ^{GEN}	29,32	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Compulsory drug treatment/drug detention centres	↑	NE	C*	33,34	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Criminalisation of drug use	↑	NE	C ^{PWID}	35					↑	NE	C ^{PWID}	35												

Notes on codes used in this table

Presence or absence of effect

- ✗ This intervention does not appear to have a significant effect upon the outcome
- ↑ This outcome may be increased by the intervention
- ↓ This outcome is decreased by the intervention
- No evidence could be located of the impact of this intervention upon the outcome
- ? unclear evidence on impact of this intervention on the outcome

Level of evidence

- A Consistent conclusions across meta-analyses, high quality systematic reviews, or multiple randomised controlled trials
- B Evidence from one or two randomised controlled trials only
- C High quality systematic reviews of cohort, case-control or cross-sectional studies
- D Systematic reviews with inconsistent conclusions from authors or cross-sectional; OR multiple consistent ecological studies
- E Cross-sectional association, case series suggesting outcome, single cohort study
- GEN Evidence drawn from people who may not specifically use any drugs (including opioids)
- PWID People who inject drugs that may not include opioids
- ALC Note that this evidence pertains to people attending alcoholics anonymous
- NE no pooled quantitative estimate reported

Intervention	Overdose				Suicide				Other injuries				Overall mortality			
	Effect	Size of effect	Level	Sources	Effect	Size of effect	Level	Sources	Effect	Size of effect	Level	Sources	Effect	Size of effect	Level	Sources
Provision of sterile injecting equipment	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Condom provision	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Naloxone	↓	-	D	36	-	-	-	-	-	-	-	-	-	-	-	-
Drug consumption rooms	↓	-	D	37	-	-	-	-	-	-	-	-	-	-	-	-
Peer-based self-help groups	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Psychosocial interventions	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Opioid detoxification alone																
Oral opioid antagonists	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Extended-release opioid antagonists	↓?	-	A	15	-	-	-	-	-	-	-	-	?	-	A	15,16
Opioid agonist treatment	↓	RaRa 0.25 (0.18, 0.36) SYNTH	C	38	↓	RaRa 0.48 (0.39, 0.59)	E	39	↓	RaRa 0.40 (0.34, 0.46)	E	39	↓	RaRa 0.33 (0.28, 0.39) SYNTH	C	38
Residential rehabilitation	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
HIV testing + informing of serostatus	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
HCV testing + informing of serostatus	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
HIV treatment	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
HCV treatment	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
STI treatment	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Medication for suicide prevention	-	-	-	-	?	-	A ^{GEN}	40	-	-	-	-	-	-	-	-
Opioid prescribing limits	↓?	-	D ^{GEN}	29	-	-	-	-	-	-	-	-	-	-	-	-
Abuse-deterrent opioid formulations	?	-	D ^{GEN}	31	-	-	-	-	-	-	-	-	?	-	D	31
Prescription opioid monitoring programmes	?	-	D ^{GEN}	32,41	-	-	-	-	-	-	-	-	?	-	D	41
Compulsory drug treatment/drug detention centres	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Criminalisation of drug use	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Notes on codes used in this table

Presence or absence of effect

✕ This intervention does not appear to have a significant effect upon the outcome

↑ This outcome may be increased by the intervention

↓ This outcome is decreased by the intervention

- No evidence could be located of the impact of this intervention upon the outcome

? unclear evidence on impact of this intervention on the outcome

Level of evidence

A Consistent conclusions across meta-analyses, high quality systematic reviews, or multiple randomised controlled trials

B Evidence from one or two randomised controlled trials only

C High quality systematic reviews of cohort, case-control or cross-sectional studies

D Systematic reviews with inconsistent conclusions from authors or cross-sectional; OR multiple consistent ecological studies

E Cross-sectional association, case series suggesting outcome, single cohort study

GEN Evidence drawn from people who may not specifically use any drugs (including opioids)

PWID People who inject drugs that may not include opioids

ALC Note that this evidence pertains to people attending alcoholics anonymous

NE no pooled quantitative estimate reported

References

1. Turner KM, Hutchinson S, Vickerman P, et al. The impact of needle and syringe provision and opiate substitution therapy on the incidence of hepatitis C virus in injecting drug users: pooling of UK evidence. *Addiction* 2011; **106**(11): 1978-88.
2. Aspinall EJ, Nambiar D, Goldberg DJ, et al. Are needle and syringe programmes associated with a reduction in HIV transmission among people who inject drugs: a systematic review and meta-analysis. *International Journal of Epidemiology* 2014; **43**(1): 235-48.
3. Platt L, Minozzi S, Reed J, et al. Needle syringe programmes and opioid substitution therapy for preventing hepatitis C transmission in people who inject drugs. *Cochrane Database Syst Rev* 2017; **9**: CD012021.
4. Larney S, Peacock A, Mathers BM, Hickman M, Degenhardt L. A systematic review of injecting-related injury and disease among people who inject drugs. *Drug Alcohol Depend* 2017; **171**: 39-49.
5. Giannou FK, Tsiara CG, Nikolopoulos GK, et al. Condom effectiveness in reducing heterosexual HIV transmission: a systematic review and meta-analysis of studies on HIV serodiscordant couples. *Expert Review of Pharmacoeconomics & Outcomes Research* 2016; **16**(4): 489-99.
6. Platt L, Minozzi S, Reed J, et al. Needle and syringe programmes and opioid substitution therapy for preventing HCV transmission among people who inject drugs: findings from a Cochrane Review and meta-analysis. *Addiction* 2018; **113**(3): 545-63.
7. Milloy MJ, Wood E. Emerging role of supervised injecting facilities in human immunodeficiency virus prevention. *Addiction* 2009; **104**(4): 620-1.
8. MacArthur GJ, van Velzen E, Palmateer N, et al. Interventions to prevent HIV and hepatitis C in people who inject drugs: A review of reviews to assess evidence of effectiveness. *International Journal of Drug Policy* 2014; **25**(1): 34-52.
9. Humphreys K, Blodgett JC, Wagner TH. Estimating the efficacy of Alcoholics Anonymous without self-selection bias: an instrumental variables re-analysis of randomized clinical trials. *Alcoholism, clinical and experimental research* 2014; **38**(11): 2688-94.
10. Gilchrist G, Swan D, Widyaratna K, et al. A Systematic Review and Meta-analysis of Psychosocial Interventions to Reduce Drug and Sexual Blood Borne Virus Risk Behaviours Among People Who Inject Drugs. *AIDS & Behavior* 2017; **21**(7): 1791-811.
11. Copenhaver MM, Johnson BT, Lee IC, Harman JJ, Carey MP. Behavioral HIV risk reduction among people who inject drugs: Meta-analytic evidence of efficacy. *Journal of Substance Abuse Treatment* 2006; **31**(2): 163-71.
12. Ivan M, Rodgers C, Maher L, van Beek I. Reducing injecting-related injury and diseases in people who inject drugs: Results from a clinician-led brief intervention. *Aust Fam Physician* 2016; **45**(3): 129-33.
13. Jahagirdar D, Wright M-D. Naltrexone for Opioid Use Disorders: A Review of Clinical Effectiveness, Cost-Effectiveness, and Guidelines Ottawa: CADTH, 2017.
14. Minozzi S, Amato L, Vecchi S, Davoli M, Kirchmayer U, Verster A. Oral naltrexone maintenance treatment for opioid dependence. *Cochrane Database of Systematic Reviews* 2011; **Issue 4, Art. No.: CD001333**.
15. Jarvis BP, Holtyn AF, Subramaniam S, et al. Extended-release injectable naltrexone for opioid use disorder: A systematic review. *Addiction* 2018; **113**: 1188-209.
16. Larney S, Gowing L, Mattick RP, Farrell M, Hall W, Degenhardt L. A systematic review and meta-analysis of naltrexone implants for the treatment of opioid dependence. *Drug and Alcohol Review* 2014; **33**(2): 115-28.

17. Gowing L, Farrell MF, Bornemann R, Sullivan LE, Ali R. Oral substitution treatment of injecting opioid users for prevention of HIV infection. *Cochrane Database Syst Rev* 2011; (8): CD004145.
18. MacArthur GJ, Minozzi S, Martin N, et al. Opiate substitution treatment and HIV transmission in people who inject drugs: Systematic review and meta-analysis. *BMJ (Online)* 2012; **345 (7879) (no pagination)**(e5945).
19. Feelemyer JP, Jarlais DCD, Arasteh K, Phillips BW, Hagan H. Changes in quality of life (WHOQOL-BREF) and addiction severity index (ASI) among participants in opioid substitution treatment (OST) in low and middle income countries: an international systematic review. *Drug & Alcohol Dependence* 2014; **134**: 251-8.
20. Malivert M, Fatseas M, Denis C, Langlois E, Auriacombe M. Effectiveness of therapeutic communities: A systematic review. *European Addiction Research* 2012; **18**(1): 1-11.
21. Schlumberger M, Desenclos J, Papaevangelou G, Richardson S, Ancelle-Park R. Knowledge of HIV serostatus and preventive behaviour among European injecting drug users: Second study. *European journal of epidemiology* 1999; **15**(3): 207-15.
22. Spelman T, Morris MD, Zang G, et al. A longitudinal study of hepatitis C virus testing and infection status notification on behaviour change in people who inject drugs. *J Epidemiol Community Health* 2015; **69**(8): 745-52.
23. Kuyper L, Milloy MJ, Marshall BD, et al. Does initiation of HIV antiretroviral therapy influence patterns of syringe lending among injection drug users? *Addict Behav* 2011; **36**(5): 560-3.
24. Reddon H, Marshall BDL, Milloy MJ. Elimination of HIV transmission through novel and established prevention strategies among people who inject drugs. *The Lancet HIV* 2019; **6**(2): e128-e36.
25. Caven M, Malaguti A, Robinson E, Fletcher E, Dillon J. Impact of Hepatitis C treatment on behavioural change in relation to drug use in people who inject drugs: a systematic review. *The International journal on drug policy* in press.
26. Manhart L, Holmes K. Randomized controlled trials of individual-level, population-level, and multilevel interventions for preventing sexually transmitted infections: what has worked? *The Journal of Infectious Diseases* 2005; **191**(S1): 7-24.
27. Shahmanesh M, Patel V, Mabey D, Cowan F. Effectiveness of interventions for the prevention of HIV and other sexually transmitted infections in female sex workers in resource poor setting: a systematic review. *Tropical Medicine & International Health* 2008; **13**(5): 659-79.
28. Elwy A, Hart G, Hawkes S, Petticrew M. Effectiveness of interventions to prevent sexually transmitted infections and human immunodeficiency virus in heterosexual men: a systematic review. *Archives of Internal Medicine* 2002; **162**(16): 1818.
29. Kertesz SG, Gordon AJ. A crisis of opioids and the limits of prescription control: United States. *Addiction* 2019; **114**(1): 169-80.
30. Rauck RL. Mitigation of IV Abuse Through the Use of Abuse-Deterrent Opioid Formulations: An Overview of Current Technologies. *Pain Practice*; **0**(0).
31. New England Comparative Effectiveness Public Advisory Council. Abuse-Deterrent Formulations of Opioids: Effectiveness and Economic Impact. New England. <https://icer-review.org/material/adf-final-report/>: Institute for Clinical and Economic Review, 2017.
32. Dowell D, Zhang K, Noonan RK, Hockenberry JM. Mandatory Provider Review And Pain Clinic Laws Reduce The Amounts Of Opioids Prescribed And Overdose Death Rates. *Health affairs (Project Hope)* 2016; **35**(10): 1876-83.
33. Werb D, Kamarulzaman A, Meacham MC, et al. The effectiveness of compulsory drug treatment: A systematic review. *The International journal on drug policy* 2016; **28**: 1-9.
34. World Health Organization. Assessment of compulsory treatment of people who use drugs in Cambodia, China, Malaysia and Viet Nam: An application of selected human rights principles. Manila: World Health Organization, 2009.
35. DeBeck K, Cheng T, Montaner JS, et al. HIV and the criminalisation of drug use among people who inject drugs: a systematic review. *The Lancet HIV* 2017; **4**(8): e357-e74.

36. Mueller SR, Walley AY, Calcaterra SL, Glanz JM, Binswanger IA. A review of opioid overdose prevention and naloxone prescribing: Implications for translating community programming into clinical practice. *Substance abuse* 2015; **36**(2): 240-53.
37. Potier C, Laprevote V, Dubois-Arber F, Cottencin O, Rolland B. Supervised injection services: What has been demonstrated? A systematic literature review. *Drug and Alcohol Dependence* 2014; **145**: 48-68.
38. Sordo L, Barrio G, Bravo MJ, et al. Mortality risk during and after opioid substitution treatment: systematic review and meta-analysis of cohort studies. *BMJ* 2017; **357**: j1550.
39. Degenhardt L, Randall D, Hall W, Law M, Butler T, Burns L. Mortality among clients of a state-wide opioid pharmacotherapy program over 20 years: risk factors and lives saved. *Drug Alcohol Depend* 2009; **105**(1-2): 9-15.
40. Hawton K, Witt KG, Taylor Salisbury TL, et al. Pharmacological interventions for self-harm in adults. *Cochrane Database of Systematic Reviews* 2015; (7).
41. Fink DS, Schleimer JP, Sarvet A, et al. Association Between Prescription Drug Monitoring Programs and Nonfatal and Fatal Drug Overdoses: A Systematic Review. *Ann Intern Med* 2018; **168**(11): 783-90.

Web appendix K: Details of literature search on cost-effectiveness studies to reduce overdose

We searched Medline for cost-effectiveness studies and other economic evaluations of opioid substitution therapy (methadone or buprenorphine, including new/emerging options such as implantable/depot formulations), naltrexone (oral or extended-release formulations), or naloxone, with overdose as the outcome of interest. Searches were completed 19 February 2018.

Search terms used were:

Group 1: Intervention		Group 2: Health economics	Results
“opioid substitution therapy” or OST or methadone or buprenorphine or “opioid agonist therapy” or “opioid replacement therapy”	+	“economic evaluation” or “cost effectiveness” or “cost benefit” or costs	176

Appendix L: Details of review on retention in treatment, dose of OST and other treatment characteristics

For data on retention in treatment, we used a recent modelling study by Mukandavire and colleagues⁶⁶ as well as an additional study supplied by one of the current paper's authors.⁶⁷ The Mukandavire study identified five observational population cohorts (with one presenting retention data separately by methadone and buprenorphine at cohort commencement) with at least five years of follow-up. Data from these studies were supplied and plotted.

We systematically reviewed published and grey literature to identify data on doses of methadone and buprenorphine prescribed in routine clinical practice, use of urine drug screening in treatment monitoring, and availability of unsupervised dosing in a country. Because the aim of the review was to describe OST as it is delivered in routine clinical practice, clinical guidelines were not consulted, and clinical trials and intervention studies were excluded. We included any cross-sectional or longitudinal observational study of people receiving OST with either methadone or buprenorphine. Studies were excluded if the sample was selected by demographic or clinical characteristics.

Search terms to identify peer-reviewed literature were as follows:

Pubmed

#	Terms	Results
1	(((methadone[Title/Abstract]) OR methadone[MeSH Terms]) OR buprenorphine[Title/Abstract]) OR buprenorphine[MeSH Terms]) OR BMT[Title/Abstract]) OR MMT[Title/Abstract]	31505
2	((((((((("opioid substitution"[Title/Abstract]) OR "opiate substitution"[Title/Abstract]) OR "opioid replacement"[Title/Abstract]) OR "opiate replacement"[Title/Abstract]) OR "opioid agonist"[Title/Abstract]) OR "opiate agonist"[Title/Abstract]) OR opiate substitution treatment[MeSH Terms]) OR OST[Title/Abstract]) OR "opioid maintenance"[Title/Abstract]) OR "opioid maintenance"[Title/Abstract]	6942
3	Search 1 and search 2	2556
4	((((((((((((("opioid substitution"[Title/Abstract]) OR "opiate substitution"[Title/Abstract]) OR "opioid replacement"[Title/Abstract]) OR "opiate replacement"[Title/Abstract]) OR "opioid agonist"[Title/Abstract]) OR "opiate agonist"[Title/Abstract]) OR opiate substitution treatment[MeSH Terms]) OR OST[Title/Abstract]) OR "opioid maintenance"[Title/Abstract]) OR "opioid maintenance"[Title/Abstract])) AND (((methadone[Title/Abstract]) OR methadone[MeSH Terms]) OR buprenorphine[Title/Abstract]) OR	811915

#	Terms	Results
	buprenorphine[MeSH Terms]) OR BMT[Title/Abstract]) OR MMT[Title/Abstract])	
5	Search 3 and search 4	399
6	((cohort[Title/Abstract]) OR longitudinal[Title/Abstract]) OR cross-sectional[Title/Abstract]	1913195
7	Search 3 and search 6	929
8	Search 3 and search 4 and search 6	157
9	Search 5 or search 7	1171
10	Search 9 restricted to post-2000	1067


Embase

#	Terms	Results
1	methadone.ti,ab OR exp Methadone/ OR buprenorphine.ti,ab OR exp Buprenorphine/ OR BMT.ti,ab OR MMT.ti,ab	60181
2	("opioid substitution" OR "opiate substitution" OR "opioid replacement" OR "opiate replacement" OR "opioid agonist" OR "opiate agonist" OR "OST" OR "opioid maintenance" OR "opiate maintenance").ti,ab OR exp Opiate Substitution Therapy/	8240
3	Search 1 or search 2	65849
4	(methadone.ti,ab OR exp Methadone/ OR buprenorphine.ti,ab OR exp Buprenorphine/ OR BMT.ti,ab OR MMT.ti,ab) OR (("opioid substitution" OR "opiate substitution" OR "opioid replacement" OR "opiate replacement" OR "opioid agonist" OR "opiate agonist" OR "OST" OR "opioid maintenance" OR "opiate maintenance").ti,ab OR exp Opiate Substitution Therapy/)	1251739
5	Search 3 and search 4	4194
6	(cohort OR longitudinal OR cross-sectional).ti,ab	2284384
7	Search 3 and search 6	16303

#	Terms	Results
8	Search 3 and search 4 and search 6	894
9	Search 5 or search 7	19603
10	Search 9 restricted to post-2000	11900


To identify data from the grey literature (i.e. reports and other material that is not formally published), we used a database collated for a systematic review of coverage of HIV prevention interventions for people who inject drugs.⁶⁸ This collection of 971 documents was assembled through extensive, systematic searching of websites related to HIV, hepatitis C virus, and/or injecting drug use. The database was searched for data on OST dose, urine drug screening, and availability of unsupervised dosing.

Figure L1: Number of people in opioid agonist treatment (OAT) per 100 people who inject drugs, 2015



Source: Figure reproduced from Larney al.¹

Figure L2: Retention in OAT observed in large-scale cohorts



Note: Data extracted from Mukandavire et al.² and Hoseinie et al.³.

References

1. Larney S, Peacock A, Leung J, et al. Global, regional, and country-level coverage of interventions to prevent and manage HIV and hepatitis C among people who inject drugs: a systematic review. *The Lancet Global Health* 2017.
2. Mukandavire C, Low A, Mburu G, et al. Impact of opioid substitution therapy on the HIV prevention benefit of antiretroviral therapy for people who inject drugs. *Aids* 2017; **31**(8): 1181-90.
3. Hoseinie L, Gholami ZH, Shadloo B, Mokri A, Amin-Esmaili M, Rahimi-Movaghar A. Drop-out from a drug treatment clinic and associated reasons. *Eastern Mediterranean Health Journal* 2017; **23**: 173-81.

Appendix M: Details of mathematical modelling

Model Description

A dynamic, deterministic, single-sex model of HIV and HCV transmission among current people who inject drugs (PWID) and people who used to inject drugs (ex-PWID) was developed. The model incorporates injecting transmission of HIV and HCV, sexual transmission of HIV among PWID, and tracks individuals following injecting cessation (ex-PWID) to fully capture HIV/HCV related mortality and life years lived. The model is stratified by injecting status (current PWID, ex-PWID), HCV disease status (susceptible, previously exposed, infected – F0, infected - F1, infected - infected - F2, infected - F3, infected - F4, infected - decompensated cirrhosis, infected - hepatocellular carcinoma), HIV disease status and stage (uninfected, acute infection, latent infection, AIDS, latent infection and on ART, AIDS and on ART), incarceration status (never incarcerated, currently incarcerated, released in the last 6 months, previously incarcerated but released longer than 6 months ago), and harm reduction status (off OAT, on OAT). Model schematics are shown in figure M1; model equations are at the end of this Appendix.

The model is open, such that individuals continually enter through initiation of injecting drug use as HCV and HIV uninfected PWID and off OAT. A proportion of PWID enter the model as never incarcerated and the remainder enter as previously incarcerated. PWID permanently cease injecting at a constant rate, whereby they leave the PWID compartment and enter the ex-PWID compartment. Individuals exit the model through mortality; PWID experience mortality through drug overdose, injury and suicide; both PWID and ex-PWID can experience HIV and HCV mortality and mortality through other causes (i.e. causes not related to suicide, drug overdose, HIV or HCV).

PWID enrol onto OAT at a time varying rate, which differs by incarceration state, and leave OAT at constant rates. A proportion of PWID entering or leaving OAT die due to the excess risk of overdose in the first 4 weeks after OAT initiation and discontinuation⁶⁹. While on OAT, PWID have a reduced risk of mortality through all causes, with different reductions in risk during periods of incarceration compared to in the community (i.e. not incarcerated)⁶⁹⁻⁷¹. While on OAT, PWID are assumed to have a reduced HIV and HCV transmission risk through injecting compared to those not on OAT^{72,73}. PWID on OAT also have greater levels of viral

suppression whilst on ART, a greater rate of ART initiation and lower rate of loss to follow up than PWID not on OAT⁷⁴.

PWID are incarcerated or re-incarcerated at constant but different rates and are released from prison at a constant rate. The rate of incarceration or reincarceration is reduced if PWID are on OAT⁷⁵. During incarceration, PWID have different overdose and suicide mortality rates compared to PWID in the community.⁷⁰ HIV and HCV transmission risk also differs in prison, and is higher in the 6 months following release than for never incarcerated PWID⁷⁶. Depending upon OAT status at time of release^{77,78}, a proportion of PWID released from prison die due to the excess risk of overdose in the first 4 weeks following prison release⁷⁹.


All PWID can acquire and transmit HCV through the sharing of injecting equipment in their given setting (prison or community). PWID susceptible to HCV become infected at a rate proportional to the chronic prevalence in their setting and the HCV transmission rate. PWID on OAT have lower HCV transmission risk, whilst recently released PWID have higher HCV transmission risk. The risk of HCV transmission may be higher or lower than in the community. PWID who are co-infected with HCV and HIV are assumed to be more likely to transmit HCV than those with HCV mono-infection. A proportion of those HCV infected (which depends upon HIV status) spontaneously clear their infection and remain in the susceptible compartment, the remainder enter the chronic HCV compartment.


Individuals with chronic HCV infection advance from F0 through F1, F2, F3, to F4 (compensated cirrhosis) according to figure 1. Rates of progression from F0 to compensated cirrhosis are elevated if individuals are HIV infected, with these rates being lowered (compared to HIV infected individuals not on ART) if they are on ART⁸⁰. Individuals with compensated cirrhosis develop decompensated cirrhosis and hepatocellular carcinoma (HCC), with individuals also developing HCC if they have decompensated cirrhosis. Individuals with decompensated cirrhosis or HCC leave the model through HCV-related mortality with the mortality rate for decompensated cirrhosis being elevated if HIV co-infected^{81,82}. No HCV treatment is assumed in this modelling because treatment rates are currently very low in these settings.

PWID susceptible to HIV are infected with a sexual and injecting force of infection which depends upon: the sexual and injecting HIV transmission rates, the proportions of PWID in


each stage of HIV infection in their setting (prison or the community), and the infectivity of each stage of infection (including ART status) which is measured relative to the latent phase of HIV infection. The model did not stratify PWID by sex and so sexual HIV transmission among PWID is modelled simply with no heterogeneity and all PWID in the community being able to mix freely to form sexual contacts. PWID on OAT have lower HIV transmission risk, whilst recently released PWID have higher HIV transmission risk through injecting. The risk of HIV transmission through injecting in prison may be higher or lower than in the community. We assume no sexual transmission whilst incarcerated due to the low levels of sex between men in prisons reported in the settings^{83,84}. Following HIV infection, individuals enter a short acute phase of HIV infection, where they are assumed to be more infectious than in the subsequent latent phase of HIV infection⁸⁵ and progress from acute HIV infection to latent HIV infection at a fixed rate. Individuals progress from latent infection to AIDS at a constant rate, where it is assumed that individuals with AIDS who are not receiving ART do not engage in injecting drug use or sexual activity and so do not contribute to HIV transmission. Conversely, individuals with AIDS who are receiving ART are assumed to have the same HIV infectiousness as individuals with latent infection who are also receiving ART. Individuals with AIDS die from HIV related-disease at a constant rate and are removed from the model. Individuals with latent HIV infection and AIDS are enrolled onto ART at a constant rate. ART enrolment rates differ for current PWID and ex-PWID and are increased if PWID are on OAT⁷⁴. Whilst on ART, individuals progress through the stages of HIV infection slower than individuals not on ART and experience a lower HIV mortality rate (when in the AIDS compartment only). The effectiveness of ART on reducing HIV infectiousness, progression and mortality depends on the proportion of individuals who are virally suppressed, which is higher amongst those receiving OAT⁷⁴. Individuals receiving ART are lost to follow up at a constant rate which depends upon their injecting status, with PWID on OAT having better ART retention than PWID not receiving OAT⁷⁴. The model assumes that these individuals can be re-enrolled onto ART at the same rate as ART-naïve individuals. ART retention and lost to follow-up are assumed to be the same in prison as the community. The proportion on ART who are virally suppressed is assumed to be the same in prison and the community for those on and off OAT but differences in OAT coverage in prison and the community results in different levels of ART coverage and viral suppression in these settings.

Figure M1: Model schematics of (a) OAT component; (b) Incarceration component; (c) HIV transmission component; (d) HCV transmission and progression component.






(c)



(d)



Model Parameterisation and Calibration

For Kiev, the model is primarily parameterised using data from three surveys; the 2015 and 2017 Alliance for Public Health Integrated Bio-Behavioural Assessment (IBBA) surveys^{86,87} and the 2015 Expanding Medication-Assisted Therapy (ExMAT) bio-behavioural survey⁸⁸. For Kentucky, the model is parameterised using data from the Social Networks Among Appalachian People study (denoted throughout as 'SNAP'), a longitudinal cohort of PWID whose methods have been described previously⁸⁹. This included data on deaths and their causes for all PWID in the cohort. Based on this data, we model an increasing PWID population in Kentucky as done previously⁹⁰. For Tehran, model parameterisation was informed by data from recent IBBAAs as well as existing published data estimates, which were obtained from an exhaustive literature source of studies of PWID in these settings. Baseline and follow-up data from the standard of care arm of a recent Vanguard randomised controlled trial study⁹¹ are used to parameterise mortality rates and causes of death in Kiev.

We calibrated the model using an approximate Bayesian computation sequential Monte Carlo scheme (ABC SMC)⁹² to setting specific data on: HIV and HCV prevalence among PWID by incarceration status (community or currently incarcerated); the difference in HIV and HCV prevalence between previously incarcerated PWID and never incarcerated PWID; the difference in HCV prevalence between HIV positive PWID and HIV negative PWID; the proportion of PWID who have ever been incarcerated, or incarcerated twice or more; the proportion of PWID currently or ever on OAT or ART; all-cause mortality rates amongst PWID; the proportion of deaths among PWID that are HIV related or due to overdose, injury, suicide or other (not HIV-related, overdose, injury or suicide). During the ABC scheme, we estimate HCV and HIV transmission (injecting and sexual) rates, mortality rates of overdose, suicide, injury and other causes (excluding HIV and HCV), incarceration and re-incarceration rates and OAT and ART enrolment rates. The ABC scheme also estimates rates of HIV and HCV transmission through injecting whilst in prison based on differences in HIV and HCV prevalence among never incarcerated, currently incarcerated and previously incarcerated PWID and estimates levels of HIV sexual transmission among the community based on differences in HCV prevalence among HIV positive and HIV negative PWID.

The ABC SMC begins with 500 parameter sets sampled from prior distributions, using Latin Hypercube sampling, which are then resampled and perturbed in an iterative manner so that each time the parameter sets better fit the data, until successive iterations no longer significantly improve the goodness of the fits as defined by a 'distance function'.

For Kentucky, this distance function is given by the log likelihood of the data given the parameters with the following contributing to the likelihood:

- HCV prevalence among never incarcerated PWID (using a beta pdf)
- HCV prevalence among previously incarcerated PWID (using a beta pdf)
- The proportion of community PWID who have ever been incarcerated (using a beta pdf)

- The proportion of community PWID who have been incarcerated twice or more (using a beta pdf)
- The proportion of PWID currently on OAT (using a beta pdf)
- The mortality rate among PWID (using a gamma pdf)
- The proportion of deaths due to each cause (using a dirichlet pdf)

For Kyiv, this distance function is given by the log likelihood of the data given the parameters with the following contributing to the likelihood:

- HCV prevalence among community PWID (using a beta pdf)
- HIV prevalence among community PWID (using a beta pdf)
- The odds ratio of the HCV prevalence among community PWID who have ever been incarcerated compared to those who have never been incarcerated (using a lognormal pdf)
- The odds ratio of the HIV prevalence among community PWID who have ever been incarcerated compared to those who have never been incarcerated (using a lognormal pdf)
- The odds ratio of the HCV prevalence among HIV positive PWID in the community compared to HIV negative PWID in the community (using a lognormal pdf)
- The proportion of community PWID who have ever been incarcerated (using a beta pdf)
- The proportion of community PWID who have been incarcerated twice or more (using a beta pdf)
- The proportion of PWID currently on OAT (using a beta pdf)
- The proportion of PWID ever on OAT (using a beta pdf)
- The mortality rate among PWID (using a gamma pdf)
- The proportion of deaths due to each cause (using a dirichlet pdf)

For Tehran, this distance function outputs a vector of two elements [a,b], with both elements of the vector being improved between iterations of the ABC algorithm. This was done to allow for additional uncertainty such that OAT and ART coverages among PWID must only fall within a range which are based on either national data or estimates from other cities in Iran. The first element (a) is a measure of how far the model projections for OAT coverage among all PWID, OAT coverage among incarcerated PWID and ART coverage among HIV positive PWID are from the corresponding ranges that are informed by data. Specifically,

$$a = \sum_{i=1,2,3} f(x_i, y_i, z_i)$$

Where

- x_1, x_2 and x_3 are the model projections for OAT coverage among all PWID, OAT coverage among incarcerated PWID and the ART coverage among HIV positive PWID, respectively.
- y_1, y_2 and y_3 are the lower limits of the target calibration range for OAT coverage among all PWID, OAT coverage among incarcerated PWID and the ART coverage among HIV positive PWID, respectively.

- z_1, z_2 and z_3 are the upper limits of the target calibration range for OAT coverage among all PWID, OAT coverage among incarcerated PWID and the ART coverage among HIV positive PWID, respectively.
- $$f(x, y, z) = \begin{cases} \frac{\left(\frac{x-y+z}{2}\right)^2}{\frac{y+z}{2}} & \text{if } x < y \text{ or } x > z \\ 0 & \text{if } y \leq x \leq z \end{cases}$$

The second element (b) is the log likelihood of the data given the parameters with the following contributing to the likelihood:

- HCV prevalence among community PWID (using a beta pdf)
- HIV prevalence among community PWID (using a beta pdf)
- HCV prevalence among currently incarcerated PWID (using a beta pdf)
- HIV prevalence among currently incarcerated PWID (using a beta pdf)
- The odds ratio of the HCV prevalence among community PWID who have ever been incarcerated compared to those who have never been incarcerated (using a lognormal pdf)
- The odds ratio of the HIV prevalence among community PWID who have ever been incarcerated compared to those who have never been incarcerated (using a lognormal pdf)
- The odds ratio of the HCV prevalence among HIV positive PWID in the community compared to HIV negative PWID in the community (using a lognormal pdf)
- The proportion of community PWID who have ever been incarcerated (using a beta pdf)
- The proportion of community PWID who have been incarcerated twice or more (using a beta pdf)
- The proportion of PWID currently on OAT (using a beta pdf)
- The proportion of PWID ever on OAT (using a beta pdf)
- The mortality rate among PWID (using a gamma pdf)
- The proportion of deaths due to each cause (using a dirichlet pdf)

Table M1 summarises the parameter values used that do not vary by setting. The increases in the risk of overdose mortality in the first 4 weeks in and out of OAT and the effects of OAT on overdose mortality in the community, HIV and HCV transmission and HIV treatment outcomes are based on pooled effect estimates from systematic reviews of international evidence. Other effects of OAT, including the effect on mortality from suicide, injury and other causes (excluding overdoses, HIV and HVC) in the community, the effect on mortality from unnatural causes (injury, suicide and overdose) in prisons and the effect on incarceration rates, are parametrised using estimates from single studies. Tables of setting specific parameter values and calibration targets are in Tables M2-M7. Model fits to data on HIV and HCV prevalence data and rates of mortality are shown in Figures M2-M9.

A simplified model without ex-PWID and HCV disease progression is used for model calibration, assuming that HCV deaths will not occur in the time frame of PWID's duration of injecting. After model calibration, for each model run we sample parameters related to HCV

disease progression and mortality, the proportion of PWID that are male and the average age that PWID start injecting from setting-specific parameter ranges to calculate mortality rates among ex-PWID, using national life expectancy data⁹³.

Table M1: Parameter distributions that do not vary by setting.

Parameter	Symbol	Parameter Value	Source/ Notes
Effects of OAT: Mortality			
Relative risk of overdose mortality if on OAT outside of prison	OAT_o^c	0.25 (95%CI: 0.18-0.36)	⁶⁹ Lognormal distribution
Relative risk of suicide mortality if on OAT outside of prison	OAT_s^c	0.48 (95%CI: 0.39-0.59)	⁷¹ Lognormal distribution
Relative risk of injury mortality if on OAT outside of prison	OAT_l^c	0.40 (95%CI: 0.34-0.46)	⁷¹ Lognormal distribution
Relative risk of other causes of mortality if on OAT	OAT_a	0.86 (95%CI: 0.75-0.99)	⁷¹ Lognormal distribution
Relative risk of suicide, injury and overdose mortality if on OAT in prison	$OAT_s^p, OAT_l^p, OAT_o^p$	0.13 (95%CI: 0.05-0.35)	⁷⁰ Lognormal distribution
Relative risk of overdose in first 4 weeks after prison release if on OAT at time of release (independent of whether retained on release).	OAT_o^{rel}	0.25 (95%CI: 0.14-0.45)	⁷¹ Lognormal distribution
Relative risk of overdose in first 4 weeks of OAT vs rest of time on OAT	OAT_{start}	1.85 (95%CI: 0.93-3.66)	^{69,94} Lognormal distribution
Relative risk of overdose in first 4 weeks off OAT vs rest of time on OAT	OAT_{leave}	1.98 (95%CI: 1.24-3.15)	^{69,94} Lognormal distribution
Effects of OAT: HIV/HCV transmission			
Relative risk of HIV IDU transmission if on OAT	OAT_{HIV}	0.46 (95%CI: 0.32-0.67)	⁷² Lognormal distribution
Relative risk of HCV IDU transmission if on OAT	OAT_{HCV}	0.50 (95%CI: 0.40-0.63)	⁷³ Lognormal distribution
Effects of OAT: HIV treatment			
Relative risk of ART recruitment if on OAT	$\frac{\Omega_1}{\Omega_0}$	1.87 (95%CI: 1.50-2.33)	⁷⁴ Lognormal distribution
Relative risk of ART LTFU if on OAT	$\frac{\Gamma_1}{\Gamma_0}$	0.77 (95%CI: 0.63-0.95)	⁷⁴ Lognormal distribution
Increase in odds of viral suppression if on OAT vs off OAT	$\frac{p_s^*(1 - p_s)}{p_s(1 - p_s^*)}$	1.45 (95%CI: 1.21-1.73)	⁷⁴ Lognormal distribution
Effects of OAT: Incarceration			
Relative rate of re/incarceration if on OAT	OAT_{inc}	0.79 (95%CI: 0.70-0.89)	⁷⁵ Lognormal distribution.
HIV Disease Progression			
Average length of HIV acute stage in months	$\frac{12}{\delta_a}$	2.90 (95%CI: 1.23-6.0)	⁸⁵ Triangular distribution.
Time from seroconversion to AIDS in years	$\frac{1}{\delta_l} + \frac{1}{\delta_a}$	Median 9.4 (IQR: 5.5-10.1)	⁹⁵ Triangular distribution.
Median time to death from AIDS in months	$\frac{12}{\mu_{HIV}}$	9.2 (IQR: 2.2-23.6)	⁹⁵ Triangular distribution.
HIV transmissibility in acute stage of infection (ω_1)	$\omega_a = \frac{\omega_1}{\omega_2}$	276 (95%CI: 131-509)	⁸⁵ Lognormal distribution

HIV transmissibility in latent stage of infection (ω_2)	$\omega_a = \frac{\omega_1}{\omega_2}$	10.6 (95%CI: 7.61-13.3)	⁸⁵ Lognormal distribution
ART Parameters			
Reduction in HIV progression if on ART and virally suppressed	t_p	0.1-0.44	⁹⁶ Lognormal distribution. ⁹⁷ Lognormal distribution. N.B. In the model, the reduction in HIV progression among those on ART who are not virally suppressed is capped at 1. ^{66,97,98}
Increase in HIV progression in non-virally suppressed on ART vs virally suppressed (VL<500)	t_s	3.07 (2.43-3.89)	
Average log viral Load if not on ART	v_b	4.79 (IQR: 4.11-5.27)	Triangular distribution – International data ^{66,97,98}
Average reduction in log viral Load if on ART and not virally suppressed	Δ_u	0.81 IQR (0-2.27)	Triangular distribution – International data ⁹⁹ Lognormal distribution – International data
Factor difference in HIV transmission risk for each log increment PVL	r_t	2.45 (95%CI: 1.85-3.26)	
HIV/HCV Transmission			
Relative increase in HCV transmission rate compared to HIV transmission rate through injecting	$\frac{\lambda}{\zeta}$	4.0-10.0	¹⁰⁰ Uniform distribution
Increase in HCV infectiousness if HIV positive	η_2	1.0-7.0	¹⁰⁰ Uniform distribution ⁷⁶ Triangular distribution: mode is average of HCV and HIV effects and range and lower/upper bounds to be the minimum/maximum of 95%CI intervals for these estimates.
Increase in HIV/HCV injecting transmission risk if currently incarcerated or recently released from prison compared to other PWID in the community	η_1	1.72 (range: 1.28 – 2.34)	

Mortality

Relative risk of overdose mortality in first 1-2 weeks after prison release vs 5-12 weeks in community (R_{r1})

$$R_r = \frac{R_{r1} + R_{r2}}{2}$$

Range: 3.2-10.0

Relative risk of overdose mortality in first 3-4 weeks after prison release vs 5-12 weeks in community (R_{r2})

$$R_r = \frac{R_{r1} + R_{r2}}{2}$$

1.7 (95%CI: 1.3-2.2)

Relative risk of overdose mortality during prison vs community

$$R_o$$

0.03-0.17

Relative risk of injury mortality during prison vs community

$$R_i$$

0.03-0.3

Relative risk of mortality through other causes (not suicide, overdose, injury, HIV or HCV-related death) during prison vs community

$$R_a$$

0.07-0.28

HCV Disease progression

% infections that spontaneous clear if HIV negative

$$\alpha_-$$

26% (95%CI: 22.0-29.0)

% decrease in odds of spontaneous clearance if HIV positive

$$\frac{\alpha_+(1 - \alpha_-)}{\alpha_-(1 - \alpha_+)}$$

0.58 (95% CI: 0.38-0.88)

Annual rate of progression from F0 to F1 in HIV negative*

$$\sigma_{3,4}^0$$

0.128 (95%CI: 0.08-0.176)

Annual rate of progression from F1 to F2 in HIV negative*

$$\sigma_{4,5}^0$$

0.059 (95%CI: 0.035-0.082)

Annual rate of progression from F2 to F3 in HIV negative*

$$\sigma_{5,6}^0$$

0.079 (95%CI: 0.056-0.10)

Annual rate of progression from F3 to F4 in HIV negative*

$$\sigma_{6,7}^0$$

0.116 (95%CI: 0.07-0.161)

Relative risk of progression to cirrhosis if on ART vs HIV positive not on ART*

$$\frac{\sigma_{3,4}^4}{\sigma_{3,4}^3}$$

0.27-0.70

Relative risk of progression to cirrhosis if HIV positive*

$$\frac{\sigma_{3,4}^3}{\sigma_{3,4}^0}$$

2.489 (95%CI: 1.811-3.42)

Annual probability of progression to decompensated cirrhosis from F4*

$$1 - e^{-\sigma_{7,8}^0}$$

0.039

Annual probability of death from decompensated cirrhosis if HIV negative*

$$1 - e^{-\mu_{dc}^0}$$

0.13

N.B. different rate used for Kentucky (See Table M4)

⁷⁹ Uniform distribution. Take average of effect w1-2 and w3-4

⁷⁹ Lognormal distribution. Take average of effect w1-2 and w3-4

^{70,71} Uniform distribution

^{70,71} Uniform distribution

^{70,71} Uniform distribution

¹⁰¹ Normal Distribution.

¹⁰² Lognormal Distribution.

¹⁰³ Normal distribution

¹⁰³ Normal distribution

¹⁰³ Normal distribution

¹⁰³ Normal distribution

^{80,104} Uniform distribution

⁸⁰ Lognormal distribution

¹⁰⁵

Beta(14.6168,360.1732)

¹⁰⁵

Beta(147.03,983.97)

Relative risk of mortality from decompensated cirrhosis if HIV positive vs negative*	$\frac{\mu_{DC}^3}{\mu_{DC}^0}$	2.26 (95%CI: 1.51-3.38)	^{81,82} Lognormal distribution
Annual probability of progression to HCC from F4 or decompensated cirrhosis*	$1 - e^{-\sigma_{7,9}^0}$	0.014	¹⁰⁵ Beta(1.9326,136.1732)
Annual probability of death from HCC*	$1 - e^{-\mu_{HCC}}$	0.43	¹⁰⁵ Beta(117.1033, 155.23)

* Parameter sampled after ABC SMC routine.

Table M2: Prior parameter distributions specific to Kentucky

Parameter	Symbol	Prior	Estimates	Source
Proportion PWID who are male		Uniform (53.6-63.5)	58.6% (53.6-63.5)	SNAP cohort ⁸³
Duration of injecting (years)	$\frac{1}{\nu}$	Uniform (5-25 years)	Mean: 10.5 (95%CI: 9.7-11.2)	SNAP cohort ⁸³
Age at first injection		Uniform: (17-26)	Median: 20.5 (IQR: 17-26)	SNAP cohort ⁸³
% of PWID who start injecting prior to first incarceration	$\frac{\phi_{0,0}^{1,0}}{\phi_{0,0}^{4,0}}$	Uniform (43.5-62.7%)	48.5% (95%CI: 43.5-53.4) have age at first injection < age at first incarceration. 57.9% (95%CI: 52.9-62.7) have age at first injection <= age at first incarceration	SNAP cohort ⁸³
Average length of one incarceration (months)	$\frac{12}{\tau}$	Normal with mean 3.40 and 95%CI: 2.70-4.09	3.40 (95% CI: 2.70 – 4.09).	SNAP cohort ⁸³
Increased risk if currently or recently incarcerated	η_1	Lognormal with mean 2.80 and 95%CI: 1.36-5.77	2.80 (95%CI: 1.36-5.77)	SNAP cohort ⁸³
OAT start date		Uniform: (1990-1999)	1990-1999	

Table M3: Data for model calibration in Kentucky

Parameter	Estimates	Source/Notes
HCV antibody prevalence		
HCV Prevalence among never incarcerated PWID	43.4% (95%CI: 38.1-48.8)	SNAP cohort ⁸³
HCV Prevalence among previously incarcerated PWID	56.6% (95%CI: 51.2-61.9)	SNAP cohort ⁸³
Incarceration data		
Proportion ever incarcerated	84.7% (95%CI: 80.8-87.9)	SNAP cohort ⁸³
Proportion incarcerated twice or more	71.2% (95%CI: 66.5-75.5%)	SNAP cohort ⁸³
OAT Coverage		
Proportion of PWID currently on OAT	4.7% (95% CI: 3.8-5.8%) in 2009.	SNAP cohort ⁸³
OAT LTFU rate.	132-352/100pyrs	SNAP cohort ⁸³
Mortality		
Overall mortality rate	0.95/100py (0.64-1.42)	SNAP cohort ⁸³
% of deaths that are		SNAP cohort ⁸³ . 27/35 observed deaths had recorded causes. 7 deaths were overdose, 2 deaths were suicides, 2 deaths were injuries, 16 deaths were other medical causes.
Overdose	25.9% (95%CI: 11.1-46.3)	
Suicide	7.4% (95%CI: 0.9-24.3)	
Injury	7.4% (95%CI: 0.9-24.3)	
Other	59.3% (95%CI: 38.8-77.6)	

Figure M2: A comparison of model fits for Kentucky with HCV prevalence estimates among (a) PWID in the community, and (b) incarcerated PWID. Continuous black line shows median projections from all the model fits, with the grey shaded areas showing 95% credibility intervals. Black points and whiskers show HCV prevalence data points with 95% confidence intervals.




Figure M3: A comparison of model fits for Kentucky with mortality estimates among current PWID. Continuous black line shows median projections from all the model fits, with the grey shaded areas showing 95% credibility intervals. Black points and whiskers show HIV prevalence data points with 95% confidence intervals.




Table M4: Prior parameter distributions specific to Kiev

Parameter	Symbol	Prior	Estimates	Source
Proportion PWID who are male		Uniform (69.9-93.1%)	74.3% (95%CI: 69.9-88.3) 90.0% (95%CI: 86.4-93.1)	ExMAT 2014 ⁸² IBBA 2015 ⁸⁰
Duration of injecting (years)	$\frac{1}{v}$	Uniform (6.5-30) 0.5* lowest estimate to 2* highest estimate	Median: 13 (IQR: 9-17) Median: 14 (IQR: 8-18) Median 15 (IQR: 8-21)	IBBA 2017 ⁸¹ ExMAT 2014 ⁸² IBBA 2015 ⁸⁰ IBBA 2017 ⁸¹
Age at first injection		Uniform (16-23)	Median: 18 (IQR:16-20) Median: 19 (IQR: 17-23) Median: 19 (IQR: 16-23)	ExMAT 2014 ⁸² IBBA 2015 ⁸⁰ IBBA 2017 ⁸¹
% of PWID who start injecting prior to first incarceration	$\frac{\phi_{0,0}^{1,0}}{\phi_{0,0}^{4,0}}$	Uniform (89.8-98.4%)	92.7% (95%CI: 89.8-94.9) have age of first injection < age of first incarceration 2.9% (1.6-5.1%) have age of first injection > age of first incarceration	ExMAT 2014 ⁸²
Average length of one incarceration (months)	$\frac{12}{\tau}$	Uniform (10.0-13.0)	Range: 10.0-13.0	ExMAT 2014 ⁸² . Range from Monte Carlo simulations to account for time spent in pre-trial detention (sizos) and prisons.
ART LTFU (per 100 py)	Γ_0	Uniform (10.6 – 16.2)	LTFU among non-PWID: 9.98 per 100 py (95%CI: 9.64-11.32); RR PWID vs non-PWID: 1.26 (95%CI: 1.10-1.43)	¹⁰⁰ Eastern Europe
% Of PWID on ART virally suppressed	p_s	Lognormal with mean 68.3% and 95%CI: 46.9-84.0)	68.3% (95%CI: 46.9-84.0)	IBBA 2017 ⁸¹ Assumed not to vary over time.

Table M5: Data for model calibration in Kiev

Parameter	Estimates	Source/Notes
HIV Prevalence		
HIV prevalence among community PWID	26.6% (95%CI: 20.3-33.3)	ExMAT 2014 ⁸²
	21.7% (95%CI: 15.7-28.2)	IBBA 2015 ⁸⁰
	25.6% (95%CI: 19.7-31.8)	IBBA 2017 ⁸¹
Odds ratio: ever incarcerated vs never incarcerated PWID	2.74 (95%CI: 1.78-4.20)	ExMAT 2014 ⁸²
	1.52 (95%CI: 0.91-2.54)	IBBA 2015 ⁸⁰
	1.98 (95%CI: 1.22-3.23)	IBBA 2017 ⁸¹
HCV antibody prevalence		
HCV prevalence among community PWID	84.8% (95%CI: 78.5-90.1)	ExMAT 2014 ⁸²
	64.4% (95%CI: 59.6-69.0)	IBBA 2015 ⁸⁰
	77.3% (95%CI: 71.1-83.1)	IBBA 2017 ⁸¹
Odds ratio: ever incarcerated vs never incarcerated PWID	1.38 (95%CI: 0.77-2.46)	ExMAT 2014 ⁸²
	1.57 (95%CI: 1.00-2.48)	IBBA 2015 ⁸⁰
	2.70 (95%CI: 1.56-4.67)	IBBA 2017 ⁸¹
Odds ratio: HIV positive PWID vs HIV negative PWID	1.04 (95%CI: 0.56-1.93)	ExMAT 2014 ⁸²
	2.32 (95%CI: 1.29-4.15)	IBBA 2015 ⁸⁰
	1.32 (95%CI: 0.73-2.40)	IBBA 2017 ⁸¹
Incarceration data		
% Ever incarcerated	49.0% (95%CI: 43.0-55.1)	ExMAT 2014 ⁸²
% Incarcerated twice or more	32.1% (95%CI: 27.6-36.9)	ExMAT 2014 ⁸²
OAT Coverage		
Proportion of PWID ever on OAT	7.8% (95%CI: 4.9-10.9)	IBBA 2013
	15.7% (95%CI: 11.0-20.9)	IBBA 2015 ⁸⁰
	21.2% (95%CI: 16.0-26.8)	IBBA 2017 ⁸¹
Proportion of PWID currently on OAT	4.4% (95%CI: 2.2-7.2)	IBBA 2015 ⁸⁰
	4.5% (95%CI: 2.0-7.6)	IBBA 2017 ⁸¹
ART Coverage		
Proportion of HIV+ve that self-report ART	20.9% (95%CI: 12.7-32.5)	ExMAT 2014 ⁸²
	25.8 (95%CI: 16.6-37.0)	IBBA 2017 ⁸¹
Mortality		
Overall mortality rate (per 100 py) among PWID	5.6 (95%CI: 3.7 – 8.3)	Vanguard ⁸⁵ HIV positive PWID contribute 184.7py and HIV negative PWID contribute 275.7py. Model mortality rate is weighted to account for this during calibration.
% of deaths that are		Vanguard ⁸⁵ . 16/25 observed deaths had recorded causes. 3 deaths were HIV related, 3 deaths were overdose, 10 deaths were other medical causes. 6 deaths among HIV positive PWID were from unknown causes (2 with CD4 counts<200). Of the unclassified deaths, we assume there were 0.5 suicide deaths, 0.5 injury deaths, 2 HIV deaths and 6 deaths from other causes.
HIV related	20% (95%CI: 6.83-40.7)	
Overdose	12% (95%CI: 2.6-31.2)	
Suicide	2.0% (95%CI: 0.0-17.2)	
Injury	2.0% (95%CI: 0.0-17.2)	
Other		
	64.0% (95%CI: 42.5-82.0)	

Figure M4: A comparison of model fits for Kiev with HIV prevalence estimates among (a) PWID in the community, and (b) incarcerated PWID. Continuous black line shows median projections from all the model fits, with the grey shaded areas showing 95% credibility intervals. Black points and whiskers show HIV prevalence data points with 95% confidence intervals.




Figure M5: A comparison of model fits for Kiev with HCV prevalence estimates among (a) PWID in the community, and (b) incarcerated PWID. Continuous black line shows median projections from all the model fits, with the grey shaded areas showing 95% credibility intervals. Black points and whiskers show HCV prevalence data points with 95% confidence intervals.




Figure M6: A comparison of model fits for Kiev with mortality estimates among current PWID. Continuous black line shows median projections from all the model fits, with the grey shaded areas showing 95% credibility intervals. Black points and whiskers show HIV prevalence data points with 95% confidence intervals. Model projections are reweighted to match the observed person years of HIV positive and HIV negative PWID in the study providing the data estimate.




Table M6: Prior parameter distributions specific to Tehran

Parameter	Symbol	Prior	Estimates	Source
Proportion PWID who are male		Uniform (92.5-99.8)	95.8% (95%CI: 94.3-97.0) 96.5% (95%CI: 94.2-98.1) 94% 97.0% (95%CI: 93.6-98.9) 96.0% (95%CI: 94.5-97.2) 94.6%: (95%CI: 92.5-96.3) 99.3% (95%CI: 98.3-99.8)	¹⁰¹ 2006/2007 ¹⁰² 2008 ¹⁰³ 2003-2005 ¹⁰⁴ 2004 ¹⁰⁵ 2006/2007 2010 BBSS ¹⁰⁶ 2014 BBSS ¹⁰⁷
Duration of injecting (years)	$\frac{1}{v}$	Uniform (2.0-18.2)	Mean: 4.533 Mean: 8.4 (95%CI: 7.9-8.9) Mean 8 (95%CI: 6.9-9.1) Mean: 6.4 (95%CI: 5.8-7.0) Neighbourhood 1: mean: 4.8 (95%CI: 4.0-5.6) Neighbourhood 2: mean: 4.9 (95%CI: 4.3-5.5)	¹⁰⁸ 2001/2002 ¹⁰¹ 2006/2007 ¹⁰⁹ 2011/2012 ¹¹⁰ 2003/2004 ¹¹¹ 2005
Age at first injection		Uniform (20-29.3)	Median: 25 Median: 26 Mean: 25.7 (95%CI: 25.2-26.2) Mean: 26.7 (95%CI: 25.7-27.7) Mean 29.3 (95%CI: 28.9-29.7) Mean: 26.1 (95%CI: 25.5-26.7) Median: 25 Median 26 (IQR: 20-30)	2010 BBSS ¹⁰⁶ 2014 BBSS ¹⁰⁷ ¹⁰¹ 2006/2007 ¹¹² 2018 ¹⁰⁹ 2011/2012 ¹¹⁰ 2003/2004 ¹¹³ 2016/2017 Maryam Alavi – unpublished data 2017/2018
% of PWID who start injecting prior to first incarceration	$\frac{\phi_{0,0}^{1,0}}{\phi_{0,0}^{4,0}}$	Uniform (43.5-98.4%)		Range of values from Kiev and Kentucky
Average length of one incarceration (year)	$\frac{12}{\tau}$	Triangular (1.0,1.14,1.5)	Mean number of incarcerations: 4.43 (95%CI: 4.13-4.75) Mean years incarcerated: 5.05 (95%CI: 4.75-5.35)	¹¹⁴ 2009 - Isfahan
Annual OAT LTFU rate	κ	Normal with mean 1.71 and 95%CI: 1.54 – 1.94	Mean retention time: 7 months (95%CI: 6.2-7.8)	¹¹⁵ 2007-2011
ART LTFU (per 100 py)	Γ_0	Uniform (47.5-97.3)	47.5-97.3	¹¹⁶ 1997-2016; Kerman
% Of PWID on ART virally suppressed	p_s	Normal with mean 60 and 95%CI: 42.1-76.1	60% (95%CI: 42.1-76.1)	¹¹⁶ 1997-2016; Kerman Assumed not to vary over time.

Table M7: Data for model calibration in Tehran

Parameter	Estimates	Source/Notes
HIV Prevalence		
HIV Prevalence among community	11.4% (95%CI: 6.4-18.4)	¹¹⁷ 2008
PWID	19.4% (95%CI: 14.7-24.7)	¹⁰² 2008
	25.7% (95%CI: 17.9-34.7)	¹⁰³ 2003-2005
	24.4% (95%CI: 20.5-28.6)	¹¹⁸ 2006
	7.9% (95%CI: 3.2-15.5)	¹⁰⁸ 2001/2002
	15.2% (95%CI: 10.1-21.5)	¹¹⁹ 2003/2004
	23.2% (95%CI: 17.6-29.5)	¹²⁰ 2004
	23.3% (95%CI: 19.9-27.1)	¹²¹ 2006/2007
	10.7% (95%CI: 8.7-12.9)	¹²² 2006/2007
	18.15% (95%CI: 15.1-21.6)	2010 BBSS ¹⁰⁶
	9.41% (95%CI: 7.1-12.2)	2014 BBSS ¹⁰⁷
Odds ratio: ever incarcerated vs never incarcerated PWID	5.57 (95%CI: 1.59-19.54)	¹¹⁹ 2003/2004
	1.68 (95%CI: 0.99-2.85)	¹²³ 2006
	1.12 (95%CI: 0.67-1.89)	¹²¹ 2006/2007
	6.83 (95%CI: 2.45-19.00)	2010 BBSS ¹⁰⁶
	8.47 (95%CI: 1.15-62.3)	2014 BBSS ¹⁰⁷
HIV Prevalence among incarcerated PWID	8.3% (95%CI: 6.6-10.4)	¹²⁴ 2013-2014
	17.1% (95%CI: 13.4-21.3)	¹⁰⁸ 2001-2002
HCV antibody prevalence		
HCV Prevalence among community	50.8% (95%CI: 41.6-60.1)	¹¹⁷ 2008
PWID	36.4% (95%CI: 28.2-45.2)	¹⁰⁸ 2001/2002
	65.1% (95%CI: 59.0-70.9)	¹⁰² 2008
	52.7% (95%CI: 43.0-62.3)	¹⁰³ 2003-2005
	28.3% (95%CI: 22.0-35.4)	¹⁰⁹ 2011/2012
	52.0% (95%CI: 44.9-59.0)	¹⁰⁴ 2004
	80.0% (95%CI: 76.2-83.6)	¹²⁵ 2006
	34.5% (95%CI: 31.4-37.7)	¹⁰⁵ 2006/2007
	61.7% (95%CI: 53.8-69.1)	Maryam Alavi – unpublished data 2018/2019
	78.7% (95%CI: 73.1-82.9%)	^{121,126} 2006/2007
Odds ratio: ever incarcerated vs never incarcerated PWID	2.18 (95%CI: 1.33-3.57)	¹¹⁸ 2006
	4.29 (95%CI: 2.94-6.27)	¹⁰⁵ 2006/2007
	2.18 (95%CI: 1.14-4.16)	Maryam Alavi – unpublished data 2018/2019
	1.35 (95%CI: 0.73-2.46)	¹²⁷ 2006/2007
HCV Prevalence among incarcerated PWID	80.6% (95%CI: 76.2-84.4)	¹⁰⁸ 2011/2002
	40.1% (95%CI: 40.3-50.3)	¹²⁸ 1995
Odds ratio: HIV positive vs HIV negative PWID	10.28 (95%CI: 6.02-17.55)	¹²² 2006/2007
	2.53 (95%CI: 1.31-4.87)	¹²⁹ 2006
Incarceration data		
% Ever incarcerated	61.1% (95%CI: 53.1-68.7)	¹³⁰ 2013/2014
	70.9% (95%CI: 67.8-73.8)	¹⁰¹ 2006/2007
	72.7% (95%CI: 67.9-77.1)	¹⁰² 2008
	61.8% (95%CI: 53.7-69.4)	¹¹⁹ 2003-2004
	75.3% (95%CI: 71.1-79.2)	¹¹⁸ 2006
	83.2% (95%CI: 79.7-86.3)	¹²¹ 2006/2007

	62.9% (95%CI: 55.1-70.2)	Maryam Alavi – unpublished data 2018/2019
	80.0% (95%CI: 76.2-83.4)	2016
	80.5% (95%CI: 77.0-83.6)	2010 BBSS ¹⁰⁶
	84.30% (95%CI: 81.1-87.1)	2014 BBSS ¹⁰⁷
% Incarcerated twice or more	61.3% (95%CI: 57.2-65.2)	2010 BBSS ¹⁰⁶
	65.3% (95%CI: 61.3-69.1)	2014 BBSS ¹⁰⁷
OAT Coverage		
Proportion of PWID currently on OAT	11% (assume between 7-15%)	UNAIDS – national programme data 2017
Proportion of incarcerated PWID currently on OAT	35-45%	Based on unpublished data from Goran prison: 45% and expert opinion for Tehran: 40%
ART Coverage		
ART coverage	11% (7-19%)	⁶² 2013
Mortality		
Overall mortality rate (per 100 py) among PWID	4.1 (95%CI: 1.3-6.9)	¹³¹ 2001-2005
	4.8 (95%CI: 0-14.3)	¹³² 2007
% of all deaths among PWID that are overdose	62.5% (24.5-91.5)	¹³¹ 2003

Figure M7: A comparison of model fits for Tehran with HIV prevalence estimates among (a) PWID in the community, and (b) incarcerated PWID. Continuous black line shows median projections from all the model fits, with the grey shaded areas showing 95% credibility intervals. Black points and whiskers show HIV prevalence data points with 95% confidence intervals.




Figure M8: A comparison of model fits for Tehran with HCV prevalence estimates among (a) PWID in the community, and (b) incarcerated PWID. Continuous black line shows median projections from all the model fits, with the grey shaded areas showing 95% credibility intervals. Black points and whiskers show HCV prevalence data points with 95% confidence intervals.





Figure M9: A comparison of model fits for Tehran with mortality estimates among current PWID. Continuous black line shows median projections from all the model fits, with the grey shaded areas showing 95% credibility intervals. Black points and whiskers show HIV prevalence data points with 95% confidence intervals.



Model Equations

$X_{i,j}^{m,n}$ and $Y_{i,j}$ are the number of PWID and ex-PWID in the model, where

- superscript m denotes incarceration status ($m = 1$, never incarcerated; $m = 2$, currently incarcerated; $m = 3$, recently released from prison (in the last 6 months); $m = 4$, previously incarcerated but not in the last 6 months)
- superscript n denotes OAT status ($n = 1$, off OAT; $n = 2$, on OAT)
- subscript i denotes HIV status ($i = 1$, Susceptible to HIV; $i = 2$, Acute HIV infection; $i = 3$, Latent HIV infection without ART; $i = 4$, Latent HIV infection and on ART; $i = 5$, AIDS without ART; $i = 6$, AIDS and on ART)
- subscript j denotes HCV status ($j = 1$, Never exposed to HCV (AB and RNA negative); $j = 2$, Previously exposed to HCV (AB positive and RNA negative); $j = 3$, Infected with F0; $j = 4$, Infected with F1; $j = 5$, Infected with F2; $j = 6$, Infected with F3; $j = 7$, Infected with F4; $j = 8$, Infected with decompensated cirrhosis ; $j = 9$, Infected with hepatocellular carcinoma)

The ordinary differential equation models can be written as

$$\frac{dX_{i,j}^{m,n}}{dt} = E_{i,j}^{m,n} + M_{i,j}^{m,n} + O_{i,j}^{m,n} + I_{i,j}^{m,n} + T_{i,j}^{m,n} + D_{i,j}^{m,n} + K_{i,j}^{m,n} + H_{i,j}^{m,n}$$

$$\frac{dY_{i,j}^{m,n}}{dt} = \bar{M}_{i,j} + \bar{D}_{i,j} + \bar{H}_{i,j}$$

The terms of these differential equations represent different aspects of the model and are described below.

Inflow of PWID

$E_{i,j}^{m,n}$ denotes the inflow of new PWID term in the equations and is given by

$$E_{i,j}^{m,n} = \phi_{i,j}^{m,n} \theta(t)$$

Where,

- $\phi_{i,j}^{m,n}$ denotes the proportion of new PWID that enter into each group
- $\theta(t)$ denotes the number of new PWID entering the model at time t

Injecting cessation and mortality

These terms in the equations are concerned with cessation of injecting as well as overdose, suicide and other mortality. $M_{i,j}^{m,n}$ and $\overline{M}_{i,j}$ denotes the terms for current PWID and ex-PWID, respectively. The terms in the equations for injecting cessation and mortality for PWID and ex-PWID are given by

$$M_{i,j}^{m,1} = -(v + \mu_1 + \mu_2 + \mu_3 + \mu_4)X_{i,j}^{m,1} \quad \text{if } m = 1,3,4;$$

$$M_{i,j}^{m,2} = -(v + OAT_s^c \mu_1 + OAT_o^c \mu_2 + OAT_l^c \mu_3 + OAT_a \mu_4)X_{i,j}^{m,2} \quad \text{if } m = 1,3,4;$$

$$M_{i,j}^{2,1} = -(v + \mu_1 + R_o \mu_2 + R_i \mu_3 + R_a \mu_4)X_{i,j}^{2,1}$$

$$M_{i,j}^{2,2} = -(v + OAT_s^p \mu_1 + OAT_o^p R_o \mu_2 + OAT_l^p R_i \mu_3 + OAT_a R_a \mu_4)X_{i,j}^{2,2}$$

$$\overline{M}_{i,j} = -\mu_5 Y_{i,j} + \sum_{n,m} v X_{i,j}^{m,n}$$

Where,

- v denotes the rate of cessation of injecting
- μ_1 denotes the suicide mortality rate in community PWID off OAT
- μ_2 denotes the overdose mortality rate in community PWID off OAT
- μ_3 denotes the injury mortality rate in community PWID off OAT
- μ_4 denotes the mortality rate among PWID due to other causes (not suicide/overdose/injury/HIV/HCV) in community PWID
- μ_5 denotes the mortality rate among Ex-PWID due to other causes (not suicide, overdose, injury, HIV or HCV-related death)
- OAT_s^p and OAT_s^c denote the relative risk of suicide mortality if on OAT vs off OAT whilst in prison and the community, respectively.
- OAT_o^p and OAT_o^c denote the relative risk of overdose mortality if on OAT vs off OAT whilst in prison and the community, respectively.
- OAT_l^p and OAT_l^c denote the relative risk of injury mortality if on OAT vs off OAT whilst in prison and the community, respectively.
- OAT_a denote the relative risk of other (not suicide, overdose, injury, HIV or HCV-related death) causes of mortality if on OAT vs off OAT whilst in prison and the community, respectively.

- R_o , R_i and R_a denote the relative risk of mortality due to overdose, injury and other causes, respectively, in prison compared to the community.

N.B. Overdose mortality due to overdose upon initiating or ending OAT is given in the section on “Transitions on and off OAT”; Overdose mortality in the first few weeks after release from prison is given in the section on “Transitions between incarceration states”; HCV mortality is in the section on HCV disease progression; and, HIV mortality is in the section on HIV disease progression.

Transitions on and off OAT

These terms in the equations are concerned with movement of current PWID onto and off of OAT and are denoted by $O_{i,j}^{m,n}$ which is given by:

$$O_{i,j}^{m,1} = -\epsilon^c X_{i,j}^{m,1} + \left(1 - \mu_2(OAT_{leave} - 1) \frac{28}{365}\right) \kappa X_{i,j}^{m,2} \quad \text{if } m = 1,3,4$$

$$O_{i,j}^{m,2} = -\kappa X_{i,j}^{m,2} + \left(1 - \mu_2 OAT_o^c(OAT_{start} - 1) \frac{28}{365}\right) \epsilon^c X_{i,j}^{m,1} \quad \text{if } m = 1,3,4$$

$$O_{i,j}^{2,1} = -\epsilon^p X_{i,j}^{2,1} + \left(1 - R_o \mu_2(OAT_{leave} - 1) \frac{28}{365}\right) \kappa X_{i,j}^{2,2}$$

$$O_{i,j}^{2,2} = -\kappa X_{i,j}^{2,2} + \left(1 - R_o \mu_2 OAT_o^p(OAT_{start} - 1) \frac{28}{365}\right) \epsilon^p X_{i,j}^{2,1}$$

Where,

- ϵ^c and ϵ^p denote the OAT recruitment rate for PWID in the community and prison, respectively.
- κ denotes the OAT loss to follow-up rate for PWID
- OAT_{start} denotes the relative increase in overdose mortality in the first four weeks after starting OAT
- OAT_{leave} denotes the relative increase in overdose mortality in the first four weeks after leaving OAT

Transitions between incarceration states

These terms in the equations are concerned with movement between incarceration states and are denoted by $I_{i,j}^{m,n}$ which is given by:

$$I_{i,j}^{1,1} = -\gamma_1 X_{i,j}^{1,1}$$

$$I_{i,j}^{1,2} = -OAT_{inc} \gamma_1 X_{i,j}^{1,2}$$

$$I_{i,j}^{2,1} = \gamma_1 X_{i,j}^{1,1} + \gamma_2 (X_{i,j}^{3,1} + X_{i,j}^{4,1}) + (1 - p_{inc}) OAT_{inc} (\gamma_1 X_{i,j}^{1,2} + \gamma_2 (X_{i,j}^{3,2} + X_{i,j}^{4,2})) - \tau X_{i,j}^{2,1}$$

$$I_{i,j}^{2,2} = p_{inc} OAT_{inc} (\gamma_1 X_{i,j}^{1,2} + \gamma_2 (X_{i,j}^{3,2} + X_{i,j}^{4,2})) - \tau X_{i,j}^{2,2}$$

$$I_{i,j}^{3,1} = \left(1 - (R_r - 1) \mu_2 \frac{28}{365}\right) \tau X_{i,j}^{2,1} + (1 - p_{rel}) \left(1 - OAT_o^{rel} (R_r - 1) \mu_2 \frac{28}{365}\right) \tau X_{i,j}^{2,2} - (F + \gamma_2) X_{i,j}^{3,1}$$

$$I_{i,j}^{3,2} = p_{rel} \left(1 - OAT_o^{rel} OAT_o^C (R_r - 1) \mu_2 \frac{28}{365}\right) \tau X_{i,j}^{2,2} - (F + \gamma_2 OAT_{inc}) X_{i,j}^{3,2}$$

$$I_{i,j}^{4,1} = 2X_{i,j}^{3,1} - \gamma_2 X_{i,j}^{4,1}$$

$$I_{i,j}^{4,2} = 2X_{i,j}^{3,2} - OAT_{inc} \gamma_2 X_{i,j}^{4,2}$$

Where,

- γ_1 denotes the incarceration rate
- γ_2 denotes the re-incarceration rate
- p_{inc} denotes the proportion of PWID that are retained on OAT upon (re-)incarceration
- p_{rel} denotes the proportion of PWID that are retained on OAT upon release from prison
- R_r denotes the relative increase in overdose mortality in the first 4 weeks after prison release vs rest of time in the community
- τ denotes the rate of release from prison
- OAT_o^{rel} denotes the relative reduction in overdose mortality in the first 4 weeks after release from prison if on OAT at time of release
- OAT_{inc} denotes the relative reduction in (re-)incarceration rates if on OAT vs off OAT.
- $1/F$ is the length of the high risk period following release from prison (0.5 year)

HCV transmission

These terms in the equations are concerned with HCV transmission and are denoted by $T_{i,j}^{m,n}$, which is given by

$$T_{i,1}^{m,1} = -\beta^m X_{i,1}^{m,1} \quad \forall i$$

$$T_{i,1}^{m,2} = -OAT_{HCV} \beta^m X_{i,1}^{m,2} \quad \forall i$$

$$T_{1,2}^{m,1} = -(1 - \alpha_-) \beta^m X_{1,2}^{m,1} + \alpha_- \beta^m X_{1,1}^{m,1}$$

$$T_{i,2}^{m,1} = -(1 - \alpha_+) \beta^m X_{i,2}^{m,1} + \alpha_+ \beta^m X_{i,1}^{m,1} \quad \text{if } i \geq 2$$

$$T_{1,2}^{m,2} = -(1 - \alpha_-) OAT_{HCV} \beta^m X_{1,2}^{m,2} + \alpha_- OAT_{HCV} \beta^m X_{1,1}^{m,2}$$

$$T_{i,2}^{m,2} = -(1 - \alpha_+) OAT_{HCV} \beta^m X_{i,2}^{m,2} + \alpha_+ OAT_{HCV} \beta^m X_{i,1}^{m,2} \quad \text{if } i \geq 2$$

$$T_{1,3}^{m,1} = (1 - \alpha_-) \beta^m (X_{1,2}^{m,1} + X_{1,1}^{m,1})$$

$$T_{i,3}^{m,1} = (1 - \alpha_+) \beta^m (X_{i,2}^{m,1} + X_{i,1}^{m,1}) \quad \text{if } i \geq 2$$

$$T_{1,3}^{m,2} = (1 - \alpha_-) OAT_{HCV} \beta^m (X_{1,2}^{m,2} + X_{1,1}^{m,2})$$

$$T_{i,3}^{m,2} = (1 - \alpha_+) OAT_{HCV} \beta^m (X_{i,2}^{m,2} + X_{i,1}^{m,2}) \quad \text{if } i \geq 2$$

$$T_{i,j}^{m,n} = 0 \quad \text{if } j > 3$$

Where,

- β^m is the HCV force of infection for PWID in incarceration state m (see section below)
- OAT_{HCV} is the relative reduction in HCV transmission risk if on OAT compared to off OAT
- α_- is the proportion of HCV infections that spontaneously clear if HIV negative
- α_+ is the proportion of HCV infections that spontaneously clear if HIV positive

N.B. The model does not include a compartment for HCV acutely infected PWID because previous modelling indicates it contributes little to transmission, probably due to the small proportion of PWID that clear infection and the relatively short duration of the acute phase¹³³. No immunity is assumed

following spontaneous clearance because the evidence is uncertain¹³⁴ and previous analyses suggested that incorporation of partial immunity has negligible impact^{133,135}.

HCV force of infection

The HCV force of infection for PWID in each incarceration state is denoted by β^m and is given by

$$\beta^1 = \lambda \frac{C_-^c + C_+^c}{T^c}$$

$$\beta^2 = \eta_1 \lambda \frac{C_-^p + C_+^p}{T^p}$$

$$\beta^3 = \eta_1 \lambda \frac{C_-^c + C_+^c}{T^c}$$

$$\beta^4 = \lambda \frac{C_-^c + C_+^c}{T^c}$$

Where,

$$C_-^c = \sum_{j \geq 3} (X_{1,j}^{1,1} + \eta_1 X_{1,j}^{3,1} + X_{1,j}^{4,1} + OAT_{HCV}(X_{1,j}^{1,2} + \eta_1 X_{1,j}^{3,2} + X_{1,j}^{4,2}))$$

$$C_+^c = \eta_2 \sum_{j \geq 3} \sum_{i > 1} [X_{i,j}^{1,1} + \eta_1 X_{i,j}^{3,1} + X_{i,j}^{4,1} + OAT_{HCV}(X_{i,j}^{1,2} + \eta_1 X_{i,j}^{3,2} + X_{i,j}^{4,2})]$$

$$T^c = \sum_{i,j} [X_{i,j}^{1,1} + \eta_1 X_{i,j}^{3,1} + X_{i,j}^{4,1} + OAT_{HCV}(X_{i,j}^{1,2} + \eta_1 X_{i,j}^{3,2} + X_{i,j}^{4,2})]$$

$$C_-^p = \sum_{j \geq 3} (X_{1,j}^{2,1} + OAT_{HCV} X_{1,j}^{2,2})$$

$$C_+^p = \eta_2 \sum_{j \geq 3} \sum_{i > 1} (X_{i,j}^{2,1} + OAT_{HCV} X_{i,j}^{2,2})$$

$$T^p = \sum_{i,j} (X_{i,j}^{2,1} + OAT_{HCV} X_{i,j}^{2,2})$$

and

- λ is the HCV transmission rate for PWID who are not on OAT and are not currently or recently incarcerated
- η_1 is the relative increase in HIV and HCV injecting risk if PWID are currently or recently incarcerated
- η_2 is the relative increase in HCV transmissibility if PWID are HIV positive compared to HIV negative

HCV disease progression

These terms in the equations are concerned with HCV disease progression and are denoted by $D_{i,j}^{m,n}$ and $\bar{D}_{i,j}$ for current PWID and ex-PWID, respectively. These are given by

$$D_{i,j}^{m,n} = 0 \quad \text{if } j = 1, 2$$

$$D_{i,3}^{m,n} = -\sigma_{3,4}^i X_{i,3}^{m,n}$$

$$D_{i,4}^{m,n} = -\sigma_{4,5}^i X_{i,4}^{m,n} + \sigma_{3,4}^i X_{i,3}^{m,n}$$

$$D_{i,5}^{m,n} = -\sigma_{5,6}^i X_{i,5}^{m,n} + \sigma_{4,5}^i X_{i,4}^{m,n}$$

$$D_{i,6}^{m,n} = -\sigma_{6,7}^i X_{i,6}^{m,n} + \sigma_{5,6}^i X_{i,5}^{m,n}$$

$$D_{i,7}^{m,n} = -(\sigma_{7,8}^i + \sigma_{7,9}^i) X_{i,7}^{m,n} + \sigma_{6,7}^i X_{i,6}^{m,n}$$

$$D_{i,8}^{m,n} = -(\sigma_{8,9}^i + \mu_{DC}^i) X_{i,8}^{m,n} + \sigma_{7,8}^i X_{i,7}^{m,n}$$

$$D_{i,9}^{m,n} = -\mu_{HCC} X_{i,9}^{m,n} + \sigma_{7,9}^i X_{i,7}^{m,n} + \sigma_{8,9}^i X_{i,8}^{m,n}$$

and

$$\bar{D}_{i,j} = 0 \quad \text{if } j = 1, 2$$

$$\bar{D}_{i,3} = -\sigma_{3,4}^i Y_{i,3}$$

$$\bar{D}_{i,4} = -\sigma_{4,5}^i Y_{i,4} + \sigma_{3,4}^i Y_{i,3}$$

$$\bar{D}_{i,5} = -\sigma_{5,6}^i Y_{i,5} + \sigma_{4,5}^i Y_{i,4}$$

$$\bar{D}_{i,6} = -\sigma_{6,7}^i Y_{i,6} + \sigma_{5,6}^i Y_{i,5}$$

$$\bar{D}_{i,7} = -(\sigma_{7,8}^i + \sigma_{7,9}^i) Y_{i,7} + \sigma_{6,7}^i Y_{i,6}$$

$$\bar{D}_{i,8} = -(\sigma_{8,9}^i + \mu_{DC}^i) Y_{i,8} + \sigma_{7,8}^i Y_{i,7}$$

$$\bar{D}_{i,9} = -\mu_{HCC} Y_{i,9} + \sigma_{7,9}^i Y_{i,7} + \sigma_{8,9}^i Y_{i,8}$$

Where,

- $\sigma_{p,q}^i$ denotes the rate of progressing from HCV disease state p to HCV disease state q if in HIV state i
- μ_{DC}^i denotes the mortality rate for HCV infected individuals with decompensated cirrhosis in HIV state i
- μ_{HCC} denotes the mortality rate for HCV infected individuals with hepatocellular carcinoma (HCC).

HIV transmission

These terms in the equations are concerned with HIV transmission and are denoted by $K_{i,j}^{m,n}$, which is given by

$$K_{1,j}^{m,1} = -(\Lambda_{inj}^m + \Lambda_{sex}^m) X_{1,j}^{m,1}$$

$$K_{2,j}^{m,1} = (\Lambda_{inj}^m + \Lambda_{sex}^m) X_{1,j}^{m,1}$$

$$K_{1,j}^{m,2} = -(OAT_{HIV} \Lambda_{inj}^m + \Lambda_{sex}^m) X_{1,j}^{m,2}$$

$$K_{2,j}^{m,2} = (OAT_{HIV} \Lambda_{inj}^m + \Lambda_{sex}^m) X_{1,j}^{m,2}$$

$$K_{i,j}^{m,n} = 0 \quad \text{if } i \geq 3$$

Where,

- Λ_{inj}^m is the HIV injecting force of infection for PWID in incarceration state m (see section below)
- Λ_{sex}^m is the HIV sexual force of infection for PWID in incarceration state m (see section below)
- OAT_{HIV} is the relative reduction in HIV injecting transmission risk if on OAT compared to off OAT

HIV injecting force of infection

The HIV injecting force of infection for PWID in each incarceration state m is denoted by Λ_{inj}^m and is given by

$$\Lambda_{inj}^1 = \zeta \frac{L_1^c + L_2^c + L_3^c}{N^c}$$

$$\Lambda_{inj}^2 = \eta_1 \zeta \frac{L^p}{N^p}$$

$$\Lambda_{inj}^3 = \eta_1 \zeta \frac{L_1^c + L_2^c + L_3^c}{N^c}$$

$$\Lambda_{inj}^4 = \zeta \frac{L_1^c + L_2^c + L_3^c}{N^c}$$

Where,

$$L_1^c = \omega_a \sum_j [X_{2,j}^{1,1} + \eta_1 X_{2,j}^{3,1} + X_{2,j}^{4,1} + OAT_{HIV}(X_{2,j}^{1,2} + \eta_1 X_{2,j}^{3,2} + X_{2,j}^{4,2})]$$

$$L_2^c = \sum_j [X_{3,j}^{1,1} + \eta_1 X_{3,j}^{3,1} + X_{3,j}^{4,1} + OAT_{HIV}(X_{3,j}^{1,2} + \eta_1 X_{3,j}^{3,2} + X_{3,j}^{4,2})]$$

$$L_3^c = \Delta \sum_{i=4,6} \sum_j [A_0(X_{i,j}^{1,1} + \eta_1 X_{i,j}^{3,1} + X_{i,j}^{4,1}) + A_1 OAT_{HIV}(X_{i,j}^{1,2} + \eta_1 X_{i,j}^{3,2} + X_{i,j}^{4,2})]$$

$$N^c = \sum_{i \neq 5} \sum_j [X_{i,j}^{1,1} + \eta_1 X_{i,j}^{3,1} + X_{i,j}^{4,1} + OAT_{HIV}(X_{i,j}^{1,2} + \eta_1 X_{i,j}^{3,2} + X_{i,j}^{4,2})]$$

$$L^p = \sum_j \left(\omega_a X_{2,j}^{2,1} + X_{3,j}^{2,1} + \Delta A_0 (X_{4,j}^{2,1} + X_{6,j}^{2,1}) + OAT_{HIV} \left(\omega_a X_{2,j}^{2,2} + X_{3,j}^{2,2} + \Delta A_1 (X_{4,j}^{2,2} + X_{6,j}^{2,2}) \right) \right)$$

$$N^p = \sum_{i \neq 5} \sum_j (X_{i,j}^{2,1} + OAT_{HIV} X_{i,j}^{2,2})$$

and,

- ζ denotes the HIV transmission rate for injecting for PWID who are not on OAT and are not currently or recently incarcerated

- ω_a denotes the relative increase in HIV transmissibility if in the acute stage of infection compared to the latent stage of infection
- A_0 denotes the relative reduction in HIV transmission for sexual transmission if on ART whilst not on OAT (see section below on Effectiveness of ART)
- A_1 denotes the relative reduction in HIV transmission for sexual transmission if on ART whilst on OAT (see section below on Effectiveness of ART)
- Δ denotes the relative reduction in the effectiveness of ART for reducing HIV transmission through injecting compared to sexual transmission.

HIV sexual force of infection

The HIV sexual force of infection for PWID in each incarceration state is denoted by Λ_{sex}^m and is given by

$$\Lambda_{sex}^m = \pi \frac{\sum_{m \neq 2} \sum_j \left(\omega_a (X_{2,j}^{m,1} + X_{2,j}^{m,2}) + X_{3,j}^{m,1} + X_{3,j}^{m,2} + A_0 (X_{4,j}^{m,1} + X_{6,j}^{m,1}) + A_1 (X_{4,j}^{m,2} + X_{6,j}^{m,2}) \right)}{\sum_{m \neq 2} \sum_n \sum_j \sum_{i \neq 5} X_{i,j}^{m,n}} \quad \text{if } m = 1, 3, 4$$

$$\Lambda_{sex}^m = 0 \quad \text{if } m = 2$$

Where π denotes the sexual HIV transmission rate.

Effectiveness of ART for reducing HIV sexual transmission

If p_s denotes the proportion of PWID on ART but not OAT who are virally suppressed, and r_s denotes the odds ratio of viral suppression when on OAT compared to PWID not on OAT then the proportion of PWID on ART with viral suppression when on OAT is given by

$$p_s^* = p_s r_s / (1 + p_s (r_s - 1)).$$

To determine the decrease in HIV transmission risk among virally suppressed and unsuppressed PWID on ART, we estimated the log difference between the baseline plasma viral load ($PVL - v_b$) for PWID off ART, and PWID on ART with suppressed (v_s) PVL $\Delta_s = v_b - v_s$, or unsuppressed (v_u) PVL $\Delta_u = v_b - v_u$. Because prior studies^{93,136} suggest HIV transmission risk increases (factor r_t) for each log increase in PVL, these log differences in PVL were used to estimate the relative decrease in transmission risk among virally suppressed (e_s) and unsuppressed (e_u) PWID on ART:

$$e_s = 1/(r_t^{\Delta_s})$$

$$e_u = 1/(r_t^{\Delta_u})$$

The average reduction in HIV transmission by ART for PWID off OAT (A_0) or on OAT (A_1) is then given by

$$A_0 = p_s e_s + (1 - p_s) e_u$$

$$A_1 = p_s^* e_s + (1 - p_s^*) e_u$$

Where,

- p_s^* and p_s denote the proportion of PWID on ART who are virally suppressed if on and off OAT, respectively
- e_s and e_u denote the relative decrease in transmission risk among virally suppressed and unsuppressed PWID on ART, respectively, compared to those not on ART. e_s can take values between 0.01 and 0.21 (i.e. between a 79% and 99% reduction) and e_u can take values between 0.07 and 1.0 (i.e. between 0% and 93% reduction).

HIV disease progression

These terms in the equations are concerned with HIV disease progression and are denoted by $H_{i,j}^{m,n}$ and $\bar{H}_{i,j}$ for current PWID and ex-PWID, respectively. These are given by

$$H_{1,j}^{m,n} = 0$$

$$H_{2,j}^{m,n} = -\delta_a X_{2,j}^{m,n}$$

$$H_{3,j}^{m,n} = -(\delta_l + \Omega_n) X_{3,j}^{m,n} + \delta_a X_{2,j}^{m,n} + \Gamma_n X_{4,j}^{m,n}$$

$$H_{4,j}^{m,n} = -(Z_n \delta_l + \Gamma_n) X_{4,j}^{m,n} + \Omega_n X_{3,j}^{m,n}$$

$$H_{5,j}^{m,n} = -(\mu_{HIV} + \Omega_n) X_{5,j}^{m,n} + \delta_l X_{3,j}^{m,n} + \Gamma_n X_{6,j}^{m,n}$$

$$H_{6,j}^{m,n} = -(Z_n \mu_{HIV} + \Gamma_n) X_{6,j}^{m,n} + \Omega_n X_{5,j}^{m,n} + Z_n \delta_l X_{4,j}^{m,n}$$

and,

$$\bar{H}_{1,j} = 0$$

$$\bar{H}_{2,j} = -\delta_a Y_{2,j}$$

$$\bar{H}_{3,j} = -(\delta_l + \Omega_2)Y_{3,j} + \delta_a Y_{2,j} + \Gamma_2 Y_{4,j}$$

$$\bar{H}_{4,j} = -(Z_2 \delta_l + \Gamma_2)Y_{4,j} + \Omega_2 Y_{3,j}$$

$$\bar{H}_{5,j} = -(\mu_{HIV} + \Omega_2)Y_{5,j} + \delta_l Y_{3,j} + \Gamma_2 Y_{6,j}$$

$$\bar{H}_{6,j} = -(Z_2 \mu_{HIV} + \Gamma_2)Y_{6,j} + \Omega_2 Y_{5,j} + Z_2 \delta_l Y_{4,j}$$

where,

- δ_a is the rate of progressing from acute HIV infection to latent HIV infection.
- δ_l is the rate of progressing from latent HIV infection to AIDS if not on ART
- μ_{HIV} is the mortality rate from AIDS if not on ART
- Ω_n is the rate of enrolling onto ART if in OAT group n
- Γ_n is the loss to follow-up rate from ART if in OAT group n
- Z_n is the relative reduction in HIV progression and mortality if on ART and in OAT group n

Effectiveness of ART for reducing HIV progression

Z_1 and Z_2 denote the relative reduction in HIV progression and mortality if on ART for those off or on OAT, respectively and are given by:

$$Z_1 = (p_s + (1 - p_s)t_s)t_p$$

$$Z_2 = (p_s^* + (1 - p_s^*)t_s)t_p$$

Where,

- p_s^* and p_s denote the proportion of PWID on ART who are virally suppressed if on and off OAT, respectively (see section on 'Effectiveness of ART for reducing HIV sexual transmission')
- t_p denotes the reduction in HIV progression if on ART and virally suppressed compare to those not on ART

- t_s denotes the relative increase in HIV progression if on ART and not virally suppressed compared to those on ART who are virally suppressed.

Model Outcomes

The number of deaths due to each cause is calculated by integrating between 2020 and 2040 the terms in the model equations related to mortality due to that cause. For example, the number of HCV deaths over 2020 and 2040 is calculated as

$$\int_{2020}^{2040} \sum_m \sum_n \sum_i (\mu_{DC}^i X_{i,8}^{m,n} + \mu_{HCC} X_{i,9}^{m,n}) + \sum_i (\mu_{DC}^i Y_{i,8} + \mu_{HCC} Y_{i,9}) dt,$$

The number of total deaths that occur between 2020 and 2040 is then given by summing the number of deaths that occur over 2020-2040 due to each cause.

The percentage of deaths averted by a scenario with OAT compared to a scenario without OAT is given by:

$$\frac{D_{No\ OAT} - D_{OAT}}{D_{No\ OAT}} * 100$$

Where,

$D_{No\ OAT}$ is the number of deaths that occur in a scenario where there is no OAT from 2020 onwards.

D_{OAT} is the number of deaths that occur in a scenario with OAT.

Finally, Life-years gained per 1000 PWID by a scenario with OAT compared to a scenario without OAT is given by:

$$\frac{L_{OAT} - L_{No\ OAT}}{N_{2020}} * 1000$$

Where,

N_{2020} is the number of PWID in the model at the start of 2020.

L_{OAT} the total number of life years in the model between 2020 and 2040 for a scenario with OAT

$L_{No\ OAT}$ the total number of life years in the model between 2020 and 2040 for a scenario where there is no OAT from 2020 onwards.

The number of life-years in the model between 2020 and 2040 is calculated by integrating the modelled population size between 2020 and 2040, i.e.

$$\int_{2020}^{2040} \sum_m \sum_n \sum_i \sum_j X_{i,j}^{m,n} + \sum_i \sum_j Y_{i,j} dt$$

It is important to note that making model projections over 20 years involves assuming that incarceration dynamics and risk behaviours associated with injecting drug use remain the same and that interventions that could impact mortality (such as naloxone) or disease transmission (such as ART) are not introduced or scaled-up. Additional model projections suggest that the impact of scaling-up OAT on mortality reduces by at most 20% (in Kiev) if we consider a shorter time frame of 10 years. The primary reason why the impact is lowered over 10 years is that the impact of scaling-up OAT on HIV deaths is significantly

reduced (by an average of 50% in Kiev) due to there being less time to accrue reductions in HIV deaths as a consequence of reductions in HIV transmission (which was the most important effect for reducing mortality in Kiev).

References

1. Degenhardt L, Bucello C, Mathers B, et al. Mortality among regular or dependent users of heroin and other opioids: a systematic review and meta-analysis of cohort studies. *Addiction* 2011; **106**(1): 32-51.
2. Hulse G, English D, Milne E, Holman C. The quantification of mortality resulting from the regular use of illicit opiates. *Addiction* 1999; **94**(2): 221-30.
3. World Health Organization. WHO Model List of Essential Medicines. World Health Organization; 2017.
4. Larance B, Degenhardt L, Lintzeris N, Winstock A, Mattick R. Definitions related to the use of pharmaceutical opioids: Extramedical use, diversion, non-adherence and aberrant medication-related behaviours. *Drug and alcohol review* 2011; **30**(3): 236-45.
5. Campbell G, Nielsen S, Larance B, et al. Pharmaceutical Opioid Use and Dependence among People Living with Chronic Pain: Associations Observed within the Pain and Opioids in Treatment (POINT) Cohort. *Pain Medicine* 2015; DOI: **10.1111/pme.12773**.
6. SAMHSA. Recovery: National and Regional Resources. <https://www.samhsa.gov/sites/default/files/samhsa-recovery-5-6-14.pdf> Accessed on 24th April 2019, 2014.
7. Jones L, Othman M, Dowswell T, et al. Pain management for women in labour: an overview of systematic reviews. *Cochrane Database of Systematic Reviews*, 2012. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD009234.pub2/abstract> (accessed).
8. Basurto OX, Rigau CD, Urrútia G. Opioids for acute pancreatitis pain. *Cochrane Database of Systematic Reviews*, 2013. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD009179.pub2/abstract> (accessed).
9. Manterola C, Vial M, Moraga J, Astudillo P. Analgesia in patients with acute abdominal pain. *Cochrane Database of Systematic Reviews*, 2011. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD005660.pub3/abstract> (accessed).
10. Toms L, Derry S, Moore RA, McQuay HJ. Single dose oral paracetamol (acetaminophen) with codeine for postoperative pain in adults. *Cochrane Database of Systematic Reviews*, 2009. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD001547.pub2/abstract> (accessed).
11. Gaskell H, Derry S, Moore RA, McQuay HJ. Single dose oral oxycodone and oxycodone plus paracetamol (acetaminophen) for acute postoperative pain in adults. *Cochrane Database of Systematic Reviews*, 2009. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD002763.pub2/abstract> (accessed).
12. Dowell D, Haegerich TM, Chou R. CDC Guideline for Prescribing Opioids for Chronic Pain--United States, 2016. *Jama* 2016; **315**(15): 1624-45.
13. Wiffen PJ, Wee B, Derry S, Bell RF, Moore RA. Opioids for cancer pain - an overview of Cochrane reviews. *Cochrane Database of Systematic Reviews*, 2017. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD012592.pub2/abstract> (accessed).
14. Wiffen PJ, Wee B, Moore RA. Oral morphine for cancer pain. *Cochrane Database of Systematic Reviews*, 2016. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD003868.pub4/abstract> (accessed).
15. Hadley G, Derry S, Moore RA, Wiffen PJ. Transdermal fentanyl for cancer pain. *Cochrane Database of Systematic Reviews*, 2013. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD010270.pub2/abstract> (accessed).
16. Schmidt-Hansen M, Bennett MI, Arnold S, Bromham N, Hilgart JS. Oxycodone for cancer-related pain. *Cochrane Database of Systematic Reviews*, 2017. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD003870.pub6/abstract> (accessed).
17. Nicholson AB, Watson GR, Derry S, Wiffen PJ. Methadone for cancer pain. *Cochrane Database of Systematic Reviews*, 2017. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD003971.pub4/abstract> (accessed).

18. Bao YJ, Hou W, Kong XY, et al. Hydromorphone for cancer pain. *Cochrane Database of Systematic Reviews*, 2016. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD011108.pub2/abstract> (accessed).
19. Wiffen PJ, Derry S, Naessens K, Bell RF. Oral tapentadol for cancer pain. *Cochrane Database of Systematic Reviews*, 2015. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD011460.pub2/abstract> (accessed).
20. Schmidt-Hansen M, Bromham N, Taubert M, Arnold S, Hilgart JS. Buprenorphine for treating cancer pain. *Cochrane Database of Systematic Reviews*, 2015. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD009596.pub4/abstract> (accessed).
21. Wiffen PJ, Derry S, Moore RA. Tramadol with or without paracetamol (acetaminophen) for cancer pain. *Cochrane Database of Systematic Reviews*, 2017. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD012508.pub2/abstract> (accessed).
22. McNicol ED, Ferguson MC, Schumann R. Methadone for neuropathic pain in adults. *Cochrane Database of Systematic Reviews*, 2017. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD012499.pub2/abstract> (accessed).
23. Haroutounian S, McNicol ED, Lipman AG. Methadone for chronic non-cancer pain in adults. *Cochrane Database of Systematic Reviews*, 2012. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD008025.pub2/abstract> (accessed).
24. Noble M, Treadwell JR, Tregear SJ, et al. Long-term opioid management for chronic noncancer pain. *Cochrane Database of Systematic Reviews*, 2010. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD006605.pub2/abstract> (accessed).
25. Gaskell H, Derry S, Stannard C, Moore RA. Oxycodone for neuropathic pain in adults. *Cochrane Database of Systematic Reviews*, 2016. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD010692.pub3/abstract> (accessed).
26. Derry S, Stannard C, Cole P, et al. Fentanyl for neuropathic pain in adults. *Cochrane Database of Systematic Reviews*, 2016. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD011605.pub2/abstract> (accessed).
27. Stannard C, Gaskell H, Derry S, et al. Hydromorphone for neuropathic pain in adults. *Cochrane Database of Systematic Reviews*, 2016. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD011604.pub2/abstract> (accessed).
28. Wiffen PJ, Derry S, Moore RA, et al. Buprenorphine for neuropathic pain in adults. *Cochrane Database of Systematic Reviews*, 2015. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD011603.pub2/abstract> (accessed).
29. Cooper TE, Fisher E, Gray AL, et al. Opioids for chronic non-cancer pain in children and adolescents. *Cochrane Database of Systematic Reviews*, 2017. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD012538.pub2/abstract> (accessed).
30. Els C, Jackson TD, Hagtvedt R, et al. High-dose opioids for chronic non-cancer pain: an overview of Cochrane Reviews. *Cochrane Database of Systematic Reviews*, 2017. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD012299.pub2/abstract> (accessed).
31. Amato L, Davoli M, Minozzi S, Ferroni E, Ali R, Ferri M. Methadone at tapered doses for the management of opioid withdrawal. *Cochrane Database of Systematic Reviews*, 2013. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD003409.pub4/abstract> (accessed).
32. Gowing L, Ali R, White JM, Mbewe D. Buprenorphine for managing opioid withdrawal. *Cochrane Database of Systematic Reviews*, 2017. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD002025.pub5/abstract> (accessed).
33. Amato L, Minozzi S, Davoli M, Vecchi S. Psychosocial and pharmacological treatments versus pharmacological treatments for opioid detoxification. *Cochrane Database of Systematic Reviews*, 2011. <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD005031.pub4/abstract>
<http://onlinelibrary.wiley.com/store/10.1002/14651858.CD005031.pub4/asset/CD005031.pdf?v=1&t=hcmqmhmn&s=403f44adf7e9c56d666c8e4496a264e748a23d1d> (accessed).

34. Osborn DA, Jeffery HE, Cole MJ. Opiate treatment for opiate withdrawal in newborn infants. *Cochrane Database of Systematic Reviews*, 2010. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD002059.pub3/abstract> (accessed).
35. Faggiano F, Vigna-Taglianti F, Versino E, Lemma P. Methadone maintenance at different dosages for opioid dependence. *Cochrane Database of Systematic Reviews*, 2003. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD002208/abstract> (accessed).
36. Mattick RP, Breen C, Kimber J, Davoli M. Buprenorphine maintenance versus placebo or methadone maintenance for opioid dependence. *Cochrane Database of Systematic Reviews*, 2014. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD002207.pub4/abstract> (accessed).
37. Ferri M, Davoli M, Perucci CA. Heroin maintenance for chronic heroin-dependent individuals. *Cochrane Database of Systematic Reviews*, 2011. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD003410.pub4/abstract> (accessed).
38. UNODC. World Drug Report 2018. Vienna: United Nations <https://www.unodc.org/wdr2018/>, 2018.
39. Ellerstrand J, Ferrari A, Degenhardt L, Whiteford H, Leung J. A systematic review of the global prevalence of prescription opioid non-medical use with an estimate of prescription opioid dependence. NDARC Technical Report No. 337. Sydney: UNSW Sydney, 2018.
40. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Int J Surg* 2010; **8**(5): 336-41.
41. GBD 2017 causes of death collaborators. Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet (London, England)* 2018; **392**(10159): 1736-88.
42. GBD 2017 DALYs and HALE Collaborators. Global, regional, and national disability-adjusted life-years (DALYs) for 359 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet (London, England)* 2018; **392**(10159): 1859-922.
43. GBD 2017 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet (London, England)* 2018; **392**(10159): 1789-858.
44. GBD 2017 Mortality Collaborators. Global, regional, and national age-sex-specific mortality and life expectancy, 1950-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet (London, England)* 2018; **392**(10159): 1684-735.
45. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Annals of internal medicine* 2009; **151**(4): 264-9.
46. Flaxman AD, Vos DT, Murray CJ. An integrative metaregression framework for descriptive epidemiology. Seattle: University of Washington Press; 2015.
47. Bucello C, Degenhardt L, Calabria B, et al. What do we know about the extent of cocaine use and dependence? Results of a global systematic review. NDARC Technical Report No. 308. Sydney: National Drug and Alcohol Research Centre, University of NSW, 2010.
48. Calabria B, Degenhardt L, Nelson P, et al. What do we know about the extent of cannabis use and dependence? Results of a global systematic review. NDARC Technical Report No. 307. Sydney: National Drug and Alcohol Research Centre, University of NSW, 2010.
49. Degenhardt L, Calabria B, Nelson P, et al. What do we know about the extent of meth/amphetamine use and dependence? Results of a global systematic review. NDARC Technical Report No. 310. Sydney: National Drug and Alcohol Research Centre, University of NSW, 2010.
50. Nelson P, McLaren J, Degenhardt L, et al. What do we know about the extent of heroin and other opioid use and dependence? Results of a global systematic review. NDARC Technical Report No. 309. Sydney: National Drug and Alcohol Research Centre, University of NSW, 2010.

51. Degenhardt L, Bucello C, Calabria B, et al. What data are available on the extent of illicit drug use and dependence globally? Results of four systematic reviews. *Drug & Alcohol Dependence* 2011; **117**: 85-101.
52. Rehm J, Frick U. Valuation of health states in the US study to establish disability weights: lessons from the literature. *International journal of methods in psychiatric research* 2010; **19**(1): 18-33.
53. Salomon JA, Haagsma JA, Davis A, et al. Disability weights for the Global Burden of Disease 2013 study. *The Lancet Global Health* 2015; **3**(11): e712-e23.
54. Haagsma JA, De Noordhout CM, Polinder S, et al. Assessing disability weights based on the responses of 30,660 people from four European countries. *Population health metrics* 2015; **13**(1): 10.
55. Salomon JA, Vos T, Hogan DR, et al. Common values in assessing health outcomes from disease and injury: disability weights measurement study for the Global Burden of Disease Study 2010. *Lancet* 2012; **380**(9859): 2129-43.
56. Strang J, Groshkova T, Uchtenhagen A, et al. Heroin on trial: systematic review and meta-analysis of randomised trials of diamorphine-prescribing as treatment for refractory heroin addiction. *Br J Psychiatry* 2015; **207**(1): 5-14.
57. Strang J, Metrebian N, Lintzeris N, et al. Supervised injectable heroin or injectable methadone versus optimised oral methadone as treatment for chronic heroin addicts in England after persistent failure in orthodox treatment (RIOTT): a randomised trial. *Lancet (London, England)* 2010; **375**(9729): 1885-95.
58. Oviedo-Joekes E, Guh D, Brissette S, et al. Hydromorphone compared with diacetylmorphine for long-term opioid dependence. *JAMA psychiatry* 2016; **73**(5): 447-55.
59. Tyndall M. An emergency response to the opioid overdose crisis in Canada: a regulated opioid distribution program. *Canadian Medical Association Journal* 2018; **15**: E35-6.
60. Mukandavire C, Low A, Mburu G, et al. Impact of opioid substitution therapy on the HIV prevention benefit of antiretroviral therapy for people who inject drugs. *Aids* 2017; **31**(8): 1181-90.
61. Hoseinie L, Gholami ZH, Shadloo B, Mokri A, Amin-Esmaeili M, Rahimi-Movaghar A. Drop-out from a drug treatment clinic and associated reasons. *Eastern Mediterranean Health Journal* 2017; **23**: 173-81.
62. Larney S, Peacock A, Leung J, et al. Global, regional, and country-level coverage of interventions to prevent and manage HIV and hepatitis C among people who inject drugs: a systematic review. *The Lancet Global Health* 2017.
63. Sordo L, Barrio G, Bravo MJ, et al. Mortality risk during and after opioid substitution treatment: systematic review and meta-analysis of cohort studies. *BMJ* 2017; **357**: j1550.
64. Larney S, Gisev N, Farrell M, et al. Opioid substitution therapy as a strategy to reduce deaths in prison: retrospective cohort study. *BMJ Open* 2014; **4**(4): e004666.
65. Degenhardt L, Randall D, Hall W, Law M, Butler T, Burns L. Mortality among clients of a state-wide opioid pharmacotherapy program over 20 years: risk factors and lives saved. *Drug and alcohol dependence* 2009; **105**(1-2): 9-15.
66. MacArthur GJ, Minozzi S, Martin N, et al. Opiate substitution treatment and HIV transmission in people who inject drugs: systematic review and meta-analysis. *BMJ* 2012; **345**(oct03 3): e5945.
67. Platt L, Minozzi S, Reed J, et al. Needle syringe programmes and opioid substitution therapy for preventing hepatitis C transmission in people who inject drugs. *Cochrane Database Syst Rev* 2017; **9**: CD012021.
68. Low AJ, Mburu G, Welton NJ, et al. Impact of Opioid Substitution Therapy on Antiretroviral Therapy Outcomes: A Systematic Review and Meta-Analysis. *Clin Infect Dis* 2016.
69. Larney S, Toson B, Burns L, Dolan K. Effect of prison-based opioid substitution treatment and post-release retention in treatment on risk of re-incarceration. *Addiction* 2012; **107**(2): 372-80.
70. Stone J, Fraser H, Lim AG, et al. Incarceration history and risk of HIV and hepatitis C virus acquisition among people who inject drugs: a systematic review and meta-analysis. *Lancet Infectious Diseases* 2018; **18**(12): 1397-409.

71. Degenhardt L, Larney S, Kimber J, et al. The impact of opioid substitution therapy on mortality post-release from prison: retrospective data linkage study. *Addiction* 2014; **109**(8): 1306-17.
72. Marsden J, Stillwell G, Jones H, et al. Does exposure to opioid substitution treatment in prison reduce the risk of death after release? A national prospective observational study in England. *Addiction* 2017; **112**(8): 1408-18.
73. Merrall EL, Kariminia A, Binswanger IA, et al. Meta-analysis of drug-related deaths soon after release from prison. *Addiction* 2010; **105**(9): 1545-54.
74. Thein H-H, Yi Q, Dore GJ, Krahn MD. Natural history of hepatitis C virus infection in HIV-infected individuals and the impact of HIV in the era of highly active antiretroviral therapy: a meta-analysis. *Aids* 2008; **22**(15): 1979-91.
75. López-Diéguez M, Montes ML, Pascual-Pareja JF, et al. The natural history of liver cirrhosis in HIV–hepatitis C virus-coinfected patients. *Aids* 2011; **25**(7): 899-904.
76. Merchante N, Girón-González JA, González-Serrano M, et al. Survival and prognostic factors of HIV-infected patients with HCV-related end-stage liver disease. *Aids* 2006; **20**(1): 49-57.
77. Demchenko I, Kostyuchok M, Byelonosova N. Monitoring of awareness, behaviour and HIV prevalence among prisoners as part of second generation HIV surveillance. 2010.
78. SeyedAlinaghi SA, Kheirandish P, Karami N, et al. High prevalence of chronic hepatitis B infection among injection drug users in Iran: the need to increase vaccination of adults at risk. *Acta Med Iran* 2010; **48**(1): 58-60.
79. Hollingsworth TD, Anderson RM, Fraser C. HIV-1 transmission, by stage of infection. *J Infect Dis* 2008; **198**(5): 687-93.
80. International HIV/AIDS Alliance in Ukraine. Integrated Behavioral and Biological Assessment 2015 - National survey of PWID. Unpublished.
81. International HIV/AIDS Alliance in Ukraine. Integrated Behavioral and Biological Assessment 2017 - National survey of PWID. Unpublished.
82. Zelenev A, Shea P, Mazhnaya A, et al. Estimating HIV and HCV prevalence among people who inject drugs in 5 Ukrainian cities using stratification-based respondent driven and random sampling. *Int J Drug Policy* 2019; **67**: 91-101.
83. Young AM, Havens JR. Transition from first illicit drug use to first injection drug use among rural Appalachian drug users: a cross-sectional comparison and retrospective survival analysis. *Addiction* 2012; **107**(3): 587-96.
84. Fraser H, Vellozzi C, Hoerger TJ, et al. Scaling-up Hepatitis C Prevention and Treatment Interventions for Achieving Elimination in the United States – a Rural and Urban Comparison. *American Journal of Epidemiology* 2019.
85. Miller WC, Hoffman IF, Hanscom BS, et al. A scalable, integrated intervention to engage people who inject drugs in HIV care and medication-assisted treatment (HPTN 074): a randomised, controlled phase 3 feasibility and efficacy study. *Lancet* 2018; **392**(10149): 747-59.
86. Toni T, Welch D, Strelkowa N, Ipsen A, Stumpf MP. Approximate Bayesian computation scheme for parameter inference and model selection in dynamical systems. *J R Soc Interface* 2009; **6**(31): 187-202.
87. United Nations, Department of Economic and Social Affairs, Population Division (2017). <https://population.un.org/wpp/DataQuery/>.
88. Hickman M, Steer C, Tilling K, et al. The impact of buprenorphine and methadone on mortality: a primary care cohort study in the United Kingdom. *Addiction* 2018; **113**(8): 1461-76.
89. Morgan D, Mahe C, Mayanja B, Okongo JM, Lubega R, Whitworth JA. HIV-1 infection in rural Africa: is there a difference in median time to AIDS and survival compared with that in industrialized countries? *AIDS* 2002; **16**(4): 597-603.
90. Nosyk B, Min JE, Evans E, et al. The effects of opioid substitution treatment and highly active antiretroviral therapy on the cause-specific risk of mortality among HIV-positive people who inject drugs. *Clinical Infectious Diseases* 2015; **61**(7): 1157-65.

91. May MT, Ingle SM, Costagliola D, et al. Cohort profile: antiretroviral therapy cohort collaboration (ART-CC). *International journal of epidemiology* 2013; **43**(3): 691-702.
92. Murray M, Hogg R, Lima V, et al. The effect of injecting drug use history on disease progression and death among HIV-positive individuals initiating combination antiretroviral therapy: collaborative cohort analysis. *HIV medicine* 2012; **13**(2): 89-97.
93. Quinn TC, Wawer MJ, Sewankambo N, et al. Viral load and heterosexual transmission of human immunodeficiency virus type 1. Rakai Project Study Group. *N Engl J Med* 2000; **342**(13): 921-9.
94. Fraser H, Mukandavire C, Martin NK, et al. HIV treatment as prevention among people who inject drugs - a re-evaluation of the evidence. *Int J Epidemiol* 2017; **46**(2): 466-78.
95. Micallef JM, Kaldor JM, Dore GJ. Spontaneous viral clearance following acute hepatitis C infection: a systematic review of longitudinal studies. *J Viral Hepat* 2006; **13**(1): 34-41.
96. Grebely J, Raffa JD, Lai C, Krajden M, Conway B, Tyndall MW. Factors associated with spontaneous clearance of hepatitis C virus among illicit drug users. *Can J Gastroenterol* 2007; **21**(7): 447-51.
97. Smith DJ, Combellick J, Jordan AE, Hagan H. Hepatitis C virus (HCV) disease progression in people who inject drugs (PWID): A systematic review and meta-analysis. *Int J Drug Policy* 2015; **26**(10): 911-21.
98. Limketkai BN, Mehta SH, Sutcliffe CG, et al. Relationship of liver disease stage and antiviral therapy with liver-related events and death in adults coinfecting with HIV/HCV. *Jama* 2012; **308**(4): 370-8.
99. Shepherd J, Jones J, Hartwell D, Davidson P, Price A, Waugh N. Interferon alfa (pegylated and non-pegylated) and ribavirin for the treatment of mild chronic hepatitis C: a systematic review and economic evaluation. 2007.
100. Mocroft A, Kirk O, Aldins P, et al. Loss to follow-up in an international, multicentre observational study. *HIV Medicine* 2008; **9**(5): 261-9.
101. Amin-Esmaili M, Rahimi-Movaghar A, Gholamrezaei M, Razaghi EM. Profile of People Who Inject Drugs in Tehran, Iran. *Acta Med Iran* 2016; **54**(12): 793-805.
102. Eskandarieh S, Nikfarjam A, Tarjoman T, Nasehi A, Jafari F, Saberi-Zafarghandi MB. Descriptive Aspects of Injection Drug Users in Iran's National Harm Reduction Program by Methadone Maintenance Treatment. *Iranian Journal of Public Health* 2013; **42**(6): 588-93.
103. Lawrinson P, Ali R, Buavirat A, et al. Key findings from the WHO collaborative study on substitution therapy for opioid dependence and HIV/AIDS. *Addiction* 2008; **103**(9): 1484-92.
104. Zamani S, Ichikawa S, Nassirimanesh B, et al. Prevalence and correlates of hepatitis C virus infection among injecting drug users in Tehran. *International Journal of Drug Policy* 2007; **18**(5): 359-63.
105. Amin-Esmaili M, Rahimi-Movaghar A, Razaghi EM, Baghestani AR, Jafari S. Factors Correlated With Hepatitis C and B Virus Infections Among Injecting Drug Users in Tehran, IR Iran. *Hepatitis Monthly* 2012; **12**(1): 23-31.
106. Khajehkazemi R, Osooli M, Sajadi L, et al. HIV prevalence and risk behaviours among people who inject drugs in Iran: the 2010 National Surveillance Survey. *Sex Transm Infect* 2013; **89**(Suppl 3): iii29-iii32.
107. Haghdoost AA, Danesh A, Sharifi H, Shokoohi M, Khajehkazemi R, Mirzazadeh A, et al. HIV Bio-Behavioral Surveillance Survey (BBSS) among People Who Inject Drugs, I. R. Iran in 2014: Project Report [In Persian]. Tehran: HIV/STI Surveillance Research Center, and WHO Collaborating Center for HIV Surveillance, Kerman University of Medical Sciences, Kerman, Iran 2014.
108. Mir-Nasseri MM, Mohammadkhani A, Tavakkoli H, Ansari E, Poustchi H. Incarceration is a major risk factor for blood-borne infection among intravenous drug users: Incarceration and blood borne infection among intravenous drug users. *Hepatitis monthly* 2011; **11**(1): 19-22.

109. Alam-Mehrjerdi Z, Moradi A, Xu F, Zarghami M, Salehi-Fadardi J, Dolan K. Willingness to Receive Treatment for Hepatitis C among Injecting Drug Users on Methadone Program: Implications for Education and Treatment. *Addict* 2016; **8**(2): 90-7.
110. Zamani S, Ono-Kihara M, Ichikawa S, Kihara M. Potential for sexual transmission of HIV infection from male injecting-drug users who have sex with men in Tehran, Iran. *Sexually Transmitted Diseases* 2010; **37**(11): 715-8.
111. Zamani S, Vazirian M, Nassirimanesh B, et al. Needle and syringe sharing practices among injecting drug users in Tehran: a comparison of two neighborhoods, one with and one without a needle and syringe program. *AIDS & Behavior* 2010; **14**(4): 885-90.
112. Noroozi M, Marshall BDL, Noroozi A, et al. Effect of Alcohol Use on Injection and Sexual Behavior among People Who Inject Drugs in Tehran, Iran: A Coarsened Exact Matching Approach. *J* 2018; **18**(2): e00416.
113. Noroozi M, Sharifi H, Noroozi A, Rezaei F, Bazrafshan MR, Armoon B. Decomposing economic disparities in risky sexual behaviors among people who inject drugs in Tehran: Blinder-Oaxaca decomposition analysis. *Epidemiol Health* 2017; **39**: e2017049.
114. Kassaian N, Adibi P, Kafashaian A, et al. Hepatitis C Virus and Associated Risk Factors among Prison Inmates with History of Drug Injection in Isfahan, Iran. *Int J Prev Med* 2012; **3**(Suppl 1): S156-61.
115. Pashaei T, Moeeni M, Roshanaei Moghdam B, Heydari H, Turner NE, Razaghi EM. Predictors of treatment retention in a major methadone maintenance treatment program in iran: a survival analysis. *J* 2014; **14**(4): 291-5.
116. HOOSHYAR SH, RANJBAR E, SHARIFI H, et al. Evaluating the HIV continuum of care and treatment in a low prevalence city in Iran: Kerman HIV-Friendly City initiative. *Assistant Director of Monitoring and Evaluation Ministry of Health, Community Development, Gender, Elderly and Children*: 7.
117. Amiri FB, Gouya MM, Saifi M, et al. Vulnerability of homeless people in Tehran, Iran, to HIV, tuberculosis and viral hepatitis. *PLoS ONE [Electronic Resource]* 2014; **9**(6): e98742.
118. Jahani MR, Kheirandish P, Hosseini M, et al. HIV seroconversion among injection drug users in detention, Tehran, Iran. *AIDS* 2009; **23**(4): 538-40.
119. Zamani S, Kihara M, Gouya MM, et al. Prevalence of and factors associated with HIV-1 infection among drug users visiting treatment centers in Tehran, Iran. *AIDS* 2005; **19**(7): 709-16.
120. Zamani S, Kihara M, Gouya MM, et al. High prevalence of HIV infection associated with incarceration among community-based injecting drug users in Tehran, Iran. *Journal of Acquired Immune Deficiency Syndromes: JAIDS* 2006; **42**(3): 342-6.
121. Malekinejad M, Mohraz M, Razani N, et al. High HIV prevalence in a respondent-driven sampling survey of injection drug users in Tehran, Iran. *AIDS & Behavior* 2015; **19**(3): 440-9.
122. Rahimi-Movaghar A, Razaghi EM, Sahimi-Izadian E, Amin-Esmaeili M. HIV, hepatitis C virus, and hepatitis B virus co-infections among injecting drug users in Tehran, Iran. *International Journal of Infectious Diseases* 2010; **14**(1): e28-33.
123. Kheirandish P, Seyedalinaghi SA, Hosseini M, et al. Prevalence and correlates of HIV infection among male injection drug users in detention in Tehran, Iran. *Journal of Acquired Immune Deficiency Syndromes: JAIDS* 2010; **53**(2): 273-5.
124. SeyedAlinaghi S, Farhodi B, Mohraz M, et al. Prevalence and Associated Factors of HIV Infection among Male Prisoners in Tehran, Iran. *Arch Iran Med* 2017; **20**(6): 356-60.
125. Kheirandish P, SeyedAlinaghi S, Jahani M, et al. Prevalence and correlates of hepatitis C infection among male injection drug users in detention, Tehran, Iran. *J Urban Health* 2009; **86**(6): 902-8.
126. Malekinejad, M. (2008). Assessment of the use of respondent -driven sampling for studying populations most at risk of HIV infection (Order No. 3353467). Available from ProQuest Dissertations & Theses Global. (304696194). Retrieved from <https://search.proquest.com/docview/304696194?accountid=9730>.

127. Malekinejad M, Navadeh S, Lotfizadeh A, Rahimi-Movaghar A, Amin-Esmaeili M, Noroozi A. High hepatitis C virus prevalence among drug users in Iran: systematic review and meta-analysis of epidemiological evidence (2001-2012). *International Journal of Infectious Diseases* 2015; **40**: 116-30.
128. ZALI MR, Aghazadeh R, Nowroozi A, Amir-Rasouly H. Anti-HCV antibody among Iranian IV drug users: is it a serious problem? 2001.
129. Hosseini M, SeyedAlinaghi S, Kheirandish P, et al. Prevalence and correlates of co-infection with human immunodeficiency virus and hepatitis C virus in male injection drug users in Iran. *Arch Iran Med* 2010; **13**(4): 318-23.
130. Asli M, Kandelouei T, Rahimyan K, Davoodbeglou F, Vaezjalali M. Characterization of Occult Hepatitis B Infection Among Injecting Drug Users in Tehran, Iran. *Hepatitis Monthly* 2016; **16**(3): e34763.
131. Jafari S, Rahimi-Movaghar A, Craib KJ, Baharlou S, Mathias R. A follow-up study of drug users in Southern Iran. *Addiction Research & Theory* 2010; **18**(1): 59-70.
132. Rahimi-Movaghar A, Khastoo G, Razzaghi E, Saberi-Zafarghandi M, Noroozi A, Jar-Siah R. Compulsory methadone maintenance treatment of severe cases of drug addiction in a residential setting in Tehran, Iran: outcome evaluation in two and six-month follow-up. *Payesh Health Monitor* 2011; **10**(4): 505-14.
133. Vickerman P, Martin N, Turner K, Hickman M. Can needle and syringe programmes and opiate substitution therapy achieve substantial reductions in hepatitis C virus prevalence? Model projections for different epidemic settings. *Addiction* 2012; **107**(11): 1984-95.
134. Grebely J, Prins M, Hellard M, et al. Hepatitis C virus clearance, reinfection, and persistence, with insights from studies of injecting drug users: towards a vaccine. *Lancet Infect Dis* 2012; **12**(5): 408-14.
135. Martin NK, Vickerman P, Foster GR, Hutchinson SJ, Goldberg DJ, Hickman M. Can antiviral therapy for hepatitis C reduce the prevalence of HCV among injecting drug user populations? A modeling analysis of its prevention utility. *J Hepatol* 2011; **54**(6): 1137-44.
136. Hughes JP, Baeten JM, Lingappa JR, et al. Determinants of per-coital-act HIV-1 infectivity among African HIV-1-serodiscordant couples. *J Infect Dis* 2012; **205**(3): 358-65.